

IMPERIAL GARDEN DUST

DIRECTIONS

Dust underneath and top of foliage thoroughly. Apply to both plants and insects. Begin application when plants are well established or when new growth appears. Dust early in the morning or evening when air is calm. Repeat every seven to ten days or as often as necessary, whether insects are present or not to prevent infestation. Dust when air is calm.

CONTROL CHART

For control of plant diseases listed, repeat application at 7-10 day intervals; more frequently under severe disease conditions.

APPLY BEFORE PLANTS ARE INFESTED AND DAMAGED BY THE FOLLOWING INSECTS AND DISEASES:

7-Day Interval Between Last Application and Harvest. **BEANS:** Mexican Bean Beetle, Leafhopper, Leaf Beetle, 12 Spotted Beetle, Ear Worm, Lygus Bug, Blister Beetle, Rust, Anthracnose, Downy Mildew. **BEETS:** (Do not apply to beets within 14 days of harvest if treated tops are to be used for food or feed.) Cabbage Worm, Flea Beetle, Downy Mildew, Cercospora Leaf Spot. **CABBAGE:** Imported Cabbage Worm, Diamondback Moth Larvae, Thrips, Cabbage Looper, Blister Beetle, Downy Mildew, Cercospora, and Alternaria Leaf Spots. **TOMATO:** Tomato Fruit Worm, Colorado Potato Beetle, Potato Aphid, Flea Beetle, Early and Late Blight, Anthracnose, Septoria Leaf Spot. **CUCUMBERS:** Striped Cucumber Beetle. **WATER-MELON, CANTALOUPE:** Striped Cucumber Beetle. Do not use sulfur on cucumbers, melons or squash unless local experience has shown that sulfur is safe on such plants. Sulfur may burn foliage when temperatures are excessively high.

14-Day Interval Between Last Application and Harvest (controls only early and mid-season diseases on these crops. For more effective control of some diseases on these crops, a fungicide alone may have to be used closer to harvest.) **STRAWBERRY:** Strawberry Leaf Roller, Strawberry Spittle Bug, Leaf Spot, Leaf Scorch. **SPINACH:** (Remove residues by washing) Flea Beetle, Webworm, Downy Mildew, White Rust. **GRAPES:** Berry Moth, Leafhopper, Black Rot, Ripe Rot. **CARROTS:** Cercospora and Alternaria Leaf Spots.

FLOWERS

CARNATIONS, CHRYSANTHEMUMS, SNAPDRAGONS: Thrips, Leaf Spot, Anthracnose. **GLADIOLUS:** Thrips, Botrytis, Curvularia, Stemphylium. **AZALEAS, CAMELIAS:** Petal Blight. Apply dust liberally to ground and litter under the plants and into the flowers of azalea, beginning when flower buds swell and repeating at 5 to 7 day intervals through blooming season. **ROSES:** Aphids, Strawberry Rootworm Beetle, Leafhopper, Black Spot.

CAUTION

May cause irritation of nose and throat! Do not breathe dust. Keep out of reach of children and domestic pets.

Wash all vegetables before using. Avoid contamination of feed and food utensils.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 307-312

M-112B-A

Imperial

ACTIVE INGREDIENTS:

Technical Methoxychlor*	3.00%
Rotenone	0.75%
Other Cube Resins	1.50%
Zineb (Zinc Ethylene Bisdithiocarbamate)**	3.25%
Sulphur	5.00%
INERT INGREDIENTS	86.50%
Total	100.00%

*Equivalent to 2.64% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.36% other isomers and related compounds.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See other cautions on back panel.

NET CONTENTS 12 OZ.

Manufactured by

IMPERIAL INC.

HENANDOAH, IOWA 51601

ACCEPTED
JUN 16 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 307-312.