

**DIRECTIONS FOR USE  
LIVESTOCK TOXAPHENE**

For controlling Hornflies, Lice and Ticks on Beef Cattle, Hogs, Sheep and Goats, dilute with water and use either as a spray or dip.

Amount Livestock Toxaphene	Amount Water
1 gallon .....	150 gallons
1 quart .....	38 gallons
1 pint .....	19 gallons
1 cup .....	9 gallons
5 tablespoons .....	3 gallons


**CAUTION:** Do not apply to dairy animals, animals within 4 weeks of slaughter or animals under 3 months of age.

**HORN FLY CONTROL:** Make first application when first horn flies are noticed in the spring. Repeat when necessary for control. Apply 1 1/2 pints of mixture to the backs of the animals only, if using low pressure (20-60 pounds) spraying equipment. If high pressure (200-400 pounds) sprayers are used, cover entire animal, using about 1 1/2 gallons of mixture per mature beef animal.

**FOR BACKRUBBER USE:** Dilute 1 cup with 4 quarts fuel oil. Apply the diluted material to saturate the backrubber mechanism. Recharge the backrubber every 2 weeks or as necessary for control. Locate backrubber to give cattle free access to it. Do not allow calves under 3 months of age, or dairy animals, or animals within 4 weeks of slaughter, to treat themselves.

L-9967A-E

EPA Reg. No. 407-322



**NET CONTENTS**

**Imperial**  
**LIVESTOCK  
TOXAPHENE**  
(emulsifiable concentrate)

(CONTAINS 6 POUNDS TECHNICAL TOXAPHENE PER GALLON)

**ACTIVE INGREDIENTS:**

Toxaphene (Technical Chlorinated Camphene containing 67% to 69% Chlorine) .....	59.0%
Petroleum Hydrocarbons .....	28.5%
<b>INERT INGREDIENTS:</b> .....	12.5%
<b>Total</b> .....	<b>100.0%</b>

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**  
See other warnings on side panel

For Controlling Hornflies, Lice and Ticks on  
Beef Cattle, Hogs, Sheep and Goats.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

Manufactured by  
**IMPERIAL INC.**  
SHENANDOAH, IOWA 51601

**DIRECTIONS FOR USE  
LIVESTOCK TOXAPHENE**

Use only on Beef Cattle, Hogs, Sheep and Goats. Do not use on animals within 4 weeks of slaughter, or under three months of age. Do not use on dairy goats.

**LICE:** Dip or spray. Thorough coverage of the entire animal is necessary for control. Treat about November 1 and repeat in 15 days. Treat sows 30 to 45 days before farrowing to prevent pigs from being infested.

**TICKS:** Dip or spray for thorough animal coverage. Apply when ticks are first noticed and repeat as necessary for control.

**WARNING:** Do not take internally. May be fatal if swallowed. **ANTIDOTE:** Give 1 tablespoon of table salt in glass of warm water and repeat until vomit is clear. Call physician. Avoid contamination of feed and food stuffs. Do not breathe spray mist. Do not get in eyes or on skin or clothing. In case of spillage on skin, wash with soap and water. Do not use on household pets, dairy animals, or within four weeks on animals being finished for slaughter, or humans. Do not use in dairy barns or milk houses. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material or washings from spray tank. Do not reuse container. Destroy when empty. This product toxic to fish and wildlife. Apply this product only as specified on this label.

**WARRANTY:** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.