

For Beef Cattle, Swine, Horses

DRY INSECTICIDE

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER CAUTIONS BELOW

ACTIVE INGREDIENT:

TOXAPHENE: (technical chlorinated camphene
containing 67% to 69% chlorine) 2.00%

INERT INGREDIENTS: 98.00%

DIRECTIONS FOR USE

As an aid to lice control on beef cattle, swine, and horses. Apply Dry Insecticide liberally in the bedding, and on the animals. For lice control on horses repeat treatment every 15 to 20 days, and on beef cattle every 10 to 15 days. Repeat treatment of swine and bedding as necessary. Do not apply on beef cattle and swine within 28 days of slaughter. Do not apply to animals under 3 months of age. Do not use on dairy cattle.

CAUTION:

Harmful if swallowed. Keep out of reach of children. Avoid breathing of dust. Avoid skin contact. Do not get in eyes. Wash thoroughly with soap and water after using. If swallowed, induce vomiting and call a physician. Do not contaminate drinking water, food, or feed.

Do not reuse container. Destroy when empty.

This product toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.


WARRANTY: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 107-323

L-9916A-E

NET CONTENTS

LBS.



Manufactured By

IMPERIAL INC.

Shenandoah, Iowa