#### IMPERIAL

### CAPTAN MOLYSLICK

### SOYBEAN SEED PROTECTALT

ACTIVE INGREDIENTS:	
Captan *	25%
IMERT INGREDIENTS "*	755
Total	100%

\* N-[(trichloromethyl) thio]-4-cyclohexene -1, 2-dicarboximide

\*\* Contains 4.75% Molybdenum equivalent from Sodium Molybdate.
15% non-oil lubricant added.

# KEEP OUT OF REACH OF CHILDREN

### DANGER

See Back Panel for Additional Precautionary Statements

Net Contents 1 Pound . WILL TREAT 4 BUSHELS

Manufactured by IMPERIAL INC. SHENANDOAH, IA 51601

ACCEPTED

SEP 1 9 1986

Under the Federal Insecticide, Fungicide, and Rodenwicks Act, as amended, for the perticide registered under EPA Reg. No. 407-325

}

)

.

,

`

of the state of

. .

e (15) १ वृद्धिः ह

از

)

.)

0

Э

7

 $\mathbf{O}$ 

Back Panel

# PRECAUTIONARY STATEMENTS HAZARDS TO HUHANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. Wear a respirator. May cause allergic skin reactions. Wash with soap and water after use. Do not get in eyes. Wear roggles or face shield when handling. Avoid contact with skin and clothing. hence and separately launder clothing before reuse. Avoid contamination of interpretation of foodstuffs. Do not use treated seed for food, feed, or oil purposes.

### TOTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of waste. Dispose of excess treated seeds or waste by burial. Apply this product only as specified on this label.

#### DERECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Hixers, loaders, and applicators, when mixing, loading, and applying, must wear mid-forearm to elbow-length natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan; boots or overshoes; one-piece overalls which have long sleeves and long pants; face shield or goggles and a hat or other appropriate head covering.

Fieldworkers and harvesters must wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan residues. Leather or fabric gloves are not acceptable.

A dust mask must be worn when mixing and loading this product.

Clothing worn while loading, mixing, and applying this product must be laundered separately from other clothing before reuse. Clothing that may have been drenched or heavily contaminated must be disposed of in accordance with State and local regulations.

## STORAGE AND DISPOSAL

STORAGE: Store in original container, in a cool, dry, secure area, away from fertilizer, food or feed. Keep container closed. Avoid excessive heat.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Captan Holyslick is formulated for planter seed box treatment of soybean seed to control damping-off, seed and soil borne fungi, such as seedling blight, seed rots and mold. The molybdenum content aids in formation of nitrogen-fixing nousle bacteria and is most beneficial in soils that are acidic, heavily leached or soil with a high iron content.

This product is designed for addition directly into planter box to reduce handling and the non-oil lubricant helps reduce cracking and splitting of soybean seed. Recommended dosage rates provide adequate coverage and any excess is deposited with seed in furrow increasing protection.

### PLANTER BOX DOSAGE RATES

Mix decage with seed in planter box with a wooden paddle or similar device. Use Captan Molyslick at the rate of 4 oz. per bushel of seed. Plant at the rate of at least 1 bushel of soybeans to the acre to insure proper Molybdenum dosage.

PLANTER EON CAPACITY DOSAGE 1 OZ. SCOOPS 1/2 Bushel 2 Scoops

1 Eushel

4 Scoops

1 1/2 Bushels

6 Scoops

For further information consult your State Agricultural Extension Service or Experiment Station.

### **IICTICE**

Selier warrants that product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, empress or implied, extends to the use of this product contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. Fc. 407-325

H-72-84-1

EPA Est. 407-IA-1

SEN SCC. NOT-IN-