

DO NOT DILUTE


Do not allow cattle to use a backrubber that is saturated to the point of run-off with this material. Do not make accessible to dairy animals. Do not spray on livestock. Do not use on calves less than 1 month old.

Use in all types of backrubbers. In automatic oilers, refill when necessary. On burlap oilers apply at the rate of one gallon per 20 linear feet of length initially. Recharge every two weeks or as necessary for control. Do not oversaturate. These devices should be made continuously accessible, one to each 35-45 head of cattle.

NOTE: Placement of backrubbers must be where cattle normally loaf, near water troughs, shade, or salt boxes. Wrap and charge brace poles and wires to allow cattle to rub their faces and heads.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



Front Panel

IMPERIAL
DRIONER[®] INSECTICIDE

ACTIVE INGREDIENTS:

Pyrethrins	1.0%
*Piperonyl Butoxide, Technical	10.0%
Amorphous Silica Gel	40.0%

INERT INGREDIENT:

Petroleum Hydrocarbons	<u>49.0%</u>
Total	100.0%

*Equivalent to 8.0% (butylcarbityl) (6-propylpiperonyl) ether and 2.0% of related compounds.

DRIONER[®] registered trademark of FMC Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid inhalation. Do not use in commercial food preparation areas, restaurants, or in edible products areas of food processing plants.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL


STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food, or feed. Keep container closed.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Wrap container and put in trash.

This product is a finely divided powder which combines the flushing and killing powers of pyrethrins and piperonyl butoxide with the dehydrative principle of thermally attrited hydrogel. It is relatively odorless and non-staining and may be used in accordance with directions in homes, stores, hotels, offices, plants, and warehouses. When used as directed, it is highly effective in controlling these insects: ANTS, ROACHES, SILVERFISH, BED BUGS, FLEAS, LICE, TICKS and DRY WOOD TERMITES.

To control ants, cockroaches, silver fish, and fleas in animal quarters, distribute Drione at the rate of 2 ounces per 1 square foot of surface area. Apply a continuous visible film underneath and behind stoves, refrigerators and appliances. Treat on, around, and behind molding, shelving, baseboards, pipe openings, cracks, crevices and other areas where insects may hide. For severe infestations apply Drione behind



cabinets and walls, in crawl spaces and attics, at the rate of 16 ounces per 1,000 square feet of surface area. Repeat treatment as necessary.

SEWER TREATMENTS: American and Oriental cockroach infestations in sanitary sewers may be treated with approximately 0.3 lb. Drione per manhole. When treating sewers, one manhole per 200 feet of line should be treated. Treatments are made by removing the manhole cover and replacing with a disc of plywood bored to receive the usual flexible hose from portable or truck-mounted ventilating equipment. 0.3 lb. of Drione is introduced into the ventilating air, to be dispersed quickly and thoroughly through the system.

BED BUGS: Take bed apart. Dust into joints and channels. If hollow, such as square or round tubing, see that the interior of framework is well dusted. Mattress should be dusted, especially tufts, folds, and edges. Picture frame moldings and all cracks and crevices in the room should be treated.

DRY WOOD TERMITES: These termites are restricted to certain parts of the country and should not be confused with Subterranean termites. Use an electric or hand operated rotary duster to distribute Drione through an entry, (e.g. crawl hole into the attic and crawl space of the building). Rate of application is recommended at one (1) pound per 1,000 sq. ft. of floor area. All wood surfaces must be covered. All vents and louvres on the windward side to be closed.

PLEAS - DOGS AND CATS: Liberally apply powder to the animal, rubbing thoroughly to the skin. Begin with the neck and include the underbody, legs, feet, and the base of the tail. Repeat if necessary. Surfaces of kennels should be dusted at rate of one ounce per 50 square feet. Heavily infested lawns, basements and crawl spaces should be dusted at rate of 16 ounces per 1,000 square feet. Repeat as necessary.

NOTICE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 407-332

A-73-84-1

EPA Est. 407-IA-1

NET CONTENTS _____
Distributed by
IMPERIAL INC.
SHENANDOAH, IA 51601

