

GENERAL INFORMATION

Midcosan 30EC is recommended for the control of soil- and seed-borne fungi which impair the germination and seedling development of field corn (planted in Illinois, Indiana, Iowa, and Missouri), and of oats, drill-seeded rice and wheat. Midcosan 30EC when used as directed, will usually increase stands and yields of field corn, oats, drill-seeded rice, and wheat by preventing seed decay and seedling diseases.

Midcosan 30EC will also be used for the control of covered smut of barley (planted in Minnesota, North Dakota, South Dakota, Montana, and Idaho) and Helminthosporium stripe of barley (planted in Maryland, Virginia, and North Carolina); loose and covered smut of oats (planted in Wisconsin, Minnesota, North Dakota, South Dakota, Montana, and Idaho); and stinking smut or bunt of wheat (planted in Minnesota, North Dakota, South Dakota, Montana, Idaho, and Washington).

DIRECTIONS FOR USE

FIELD CORN: Seed Protection — For ready-mix treaters, use 2.0 fluid ounces of Midcosan 30EC per 100 pounds of corn seed. Midcosan 30EC may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 2.0 fluid ounces of Midcosan 30EC in 10 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater, provided that 2.0 fluid ounces of Midcosan 30EC is applied per 100 pounds of seed.

BARLEY, OATS, DRILL-SEEDED RICE, and WHEAT: Seed Protection — For ready-mix treaters, use 0.75 fluid ounce of Midcosan 30EC per bushel of seed. Midcosan 30EC may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 0.75 fluid ounce of Midcosan 30EC in 4 fluid ounces of water and apply this dispersion per bushel of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater, provided that 0.75 fluid ounce of Midcosan 30EC is applied per bushel of seed.

Imperial

MIDCOSAN 30EC

ACCEPTED

March 6 1973
UNDER THE FEDERAL INSEC
FUNGICIDE AND RODENTICID
FOR ECONOMIC POISON REG
ED UNDER NO. 340

COMPOSITION

ACTIVE INGREDIENT:

2-(thiocyanomethylthio) benzothiazole 30 percent

INERT INGREDIENTS 70 percent

DANGER

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS

Manufactured By

IMPERIAL INC.

SHENANDOAH, IOWA 51601

Imperial

ACCEPTED

March 6 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 107-340

DANGER

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. If swallowed, call a physician immediately. Give patient doses of powdered charcoal immediately or all he can swallow of raw egg whites, milk, gruel, or flour and water. Then induce vomiting with soap or mustard in water. May be harmful if absorbed through the skin. Workers handling the product should wear rubber gloves and goggles and should avoid contact of the product with clothing or skin. After exposure, wash promptly and thoroughly with an abrasive soap and cool water and finally with glycerin. In case of persistent irritation of the skin, obtain medical attention. If the product gets into the eyes, it causes eye damage; therefore, flood eyes immediately and thoroughly with cool water for 15 to 30 minutes. See a physician if any irritation persists. Keep the drum closed when not in use.

The product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment and disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

MIDCOSAN 30EC

COMPOSITION

ACTIVE INGREDIENT:
2- (thiocyanomethylthio) benzothiazole 30 percent
INERT INGREDIENTS 70 percent

DANGER

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS - - -

Manufactured By

IMPERIAL INC.

SHENANDOAH, IOWA 51601

IMPORTANT

Do not use crops grown from Midcosan 30EC treated seed for grazing of livestock or to harvest as silage or forage. However, grain harvested from plants grown from Midcosan 30EC treated seed can be used for food or feed.

Label treated seed. Seeds treated with Midcosan 30EC cannot be used for food, feed, oil production, or any other purpose except planting.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Reg. No. 107-340

M-580B-B