


Imperial



# RON-PONA FLY BAIT

CONTAINS KORLAN\* AND VAPONA<sup>R</sup>  
CONTROLS RESISTANT AND NON-RESISTANT FLIES

<b>ACTIVE INGREDIENTS:</b>	
Ronnel (O,O-Dimethyl O-(2,4,5-trichlorophenyl) phosphorothioate)	.500%
2,2-dichlorovinyl dimethyl phosphate	.225**
Related Compounds	.022**
<b>INERT INGREDIENTS:</b>	99.250%

\*\*Equivalent to 2.5% W Vapona<sup>R</sup> Insecticide  
\*Korlan<sup>R</sup> — Registered trademark Dow Chemical Co.  
Vapona<sup>R</sup> — Registered trademark Shell Chemical Co.

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating and smoking. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

See Other Cautions on Side Panel

NET WEIGHT

Manufactured by  
**IMPERIAL INC.**  
SHENANDOAH, IOWA 51601

## DIRECTIONS FOR USE FOR FLY CONTROL

**Dry Bait:** Scatter bait where flies congregate.  
**Liquid Bait:** Dissolve 2 lbs. of bait in 1 gallon of water and apply as a coarse spray or sprinkle. Place bait in areas inaccessible to children and pets.

**Dairy Barns, Milk Sheds, Livestock Barns, Loafing Sheds, Pig Pens, Stables and Outdoor Areas:** Distribute bait at the rate of 1/4 pound (1 qt. liquid bait) per 500 sq. feet. Apply to floors, window sills, walkways and manure piles where flies are active. On freshly limed surfaces, place treatment on paper. Repeat as needed for control. Do not contaminate milk or milking equipment.

**Poultry Houses:** Distribute bait at rate of 1/4 pound (1 qt. liquid bait) per 500 sq. feet where flies congregate, but keep out of reach of birds. Apply to manure, window sills, outside of penned enclosures and feed storage areas. Repeat as necessary.

**Caged Poultry and Turkeys on Wire:** Distribute bait at the rate of 1/4 pound (1 qt. liquid bait) per 500 sq. feet on droppings under cages or perches. Repeat application at weekly intervals.

**Dog Kennels:** Distribute bait at the rate of 1/4 pound (1 qt. liquid bait) per 500 sq. feet. Apply to outside of dog run ways, on window sills, ledges

and other surfaces out of reach of animals. For caged units apply to dropping area.

**For Reducing Population of Resistant and Non-Resistant House Flies In and Around Outdoor Living Areas, Picnic Grounds, Backyard Areas, Patios, Loading Docks, Outdoor Latrines, Parking Areas, Refuse Areas Around Service Stations, Open Air Drive-Ins, Ice Cream Stands, Garbage Collection and Disposal Areas:** Distribute the bait at the rate of 1/2 pound (2 qt. liquid bait) per 500 sq. feet. Repeat as necessary for control.

**Garbage Cans:** Apply the dry bait around and inside garbage cans. Repeat as necessary for control.

**Caution:** Place bait in areas inaccessible to children.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of waste. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

**Notice:** Seller makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning the use of this product or uses which extend beyond the use of the product under normal conditions in accord with the statements on this label.

EPV REG. 1007-112

M-F 10-5 P.M.