

LO-VOL SUPER BRUSH KILL

For control of mixed stands of brush consisting of sumac, elm, willow, poison ivy, and certain other softwoods as well as osage orange (hedge), hickory, oak, and other hardwoods. LO-VOL Super Brush Kill is particularly effective for basal, stump, frill and foliage treatments in non-crop areas such as waste lands, fence rows, and for spot treatment in pastures.

DO NOT graze dairy animals on treated areas within six weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

DIRECTIONS

FOLIAGE (leaf) SPRAYING

Apply LO-VOL Super Brush Kill to woody growth anytime after foliage is fully out until three weeks before frost. Best results are received from applications in late spring or early summer when leaves are fully out and before hot weather. Applications must be complete, and all parts of the plants, including foliage, stems and bark, wet with spray. Any size brush can be foliage sprayed, but basal or stump treatment will generally be more economical for brush over 8 to 10 feet high. For leaf spraying, mix only with water as shown below:


DILUTION TABLE — LEAF SPRAYING

LO-VOL Super Brush Kill	Water
2 tablespoons	1 Gallon
2½ to 3½ cups	30 Gallons
2 to 3 pints	50 Gallons
2 to 3 quarts	100 Gallons

The lower dilution rates may be used when spraying just after the leaves are out. Repeat applications may be necessary if new growth develops, but a single treatment in any one season is usually sufficient. Broad-leaved weeds are also controlled by this application.

BASAL (TRUNK) TREATMENT

Dilute LO-VOL Super Brush Kill at the rate of 1 pint to 3 gallons or 1 gallon to 25 gallons of fuel oil. Spray from the ground line up 18 inches on the trunk. Wet trunk on all sides and all exposed roots until run-off occurs. Use low pressure spray. Treat any time of year.



Imperial

TREAT:

- LEAVES
- STUMPS
- TRUNKS

LO-VOL SUPER BRUSH KILLER

CONTROLS:

- OAK
 - ELM
 - HICKORY
 - WILLOW
 - OSAGE ORANGE
 - SUMAC
 - PERSIMMON
- and certain other woody plants

A LOW VOLATILE HERBICIDE

ACTIVE INGREDIENTS:	
*Mixed Octyl Ester of 2,4-Dichlorophenoxyacetic acid (minimum)	34.43%
**Mixed Octyl Ester of 2,4,5-Trichlorophenoxyacetic acid (minimum)	32.89%
INERT INGREDIENTS	
Total	100.00%

*Contains 2 pounds 2,4-Dichlorophenoxyacetic acid per gallon, equivalent to 22.85%.
 **Contains 2 pounds 2,4,5-Trichlorophenoxyacetic acid per gallon, equivalent to 22.85%.

CAUTION:

**KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL.**

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

Manufactured by
IMPERIAL INC.
SHENANDOAH, IOWA 51601

NET CONTENTS:



STUMP TREATMENT

Dilute LO-VOL Super Brush Kill with fuel oil as for Basal Treatment. Apply as coarse spray. Wet the freshly cut surface, bark, and exposed roots down to the ground until run-off occurs. Treat immediately after cutting any time during the year.

FRILL TREATMENT

Dilute LO-VOL Super Brush Kill with fuel oil as for Basal Treatment. On trees over 6 inches in diameter or thick-barked species cut overlapping notches in a ring around the trunk near base. Cut through bark, but do not remove chips. Apply to thoroughly saturate freshly cut surfaces at any time of year.

CAUTION: HARMFUL IF SWALLOWED. Avoid contact with eyes, skin or clothing. In case of contact, wash with soap and water.

DO NOT use around homes, recreational areas, and similar sites.

Avoid spray drift to susceptible plants such as cotton, soybeans, tomatoes, grapes, sugar beets, clover, alfalfa, ornamentals, fruit trees and food crops. Coarse sprays are less likely to drift. At high temperatures vapors from this product may injure susceptible plants in the immediate vicinity—application by trunk treatment method during the dormant season will usually eliminate this hazard. Do not use the same spraying equipment for other purposes until thoroughly cleaned with a suitable chemical cleaner. Do not store near fertilizers, seeds, insecticides or fungicides.

This product is toxic to fish. Avoid contamination of irrigation ditches, lakes, streams, ponds, ditch banks or water used for domestic purposes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

APPLY ONLY AS DIRECTED ON THIS LABEL

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

NOTICE

Seller guarantee shall be limited to the terms of the label, and subject thereto, the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

LO-VOL SUPER BRUSH KILL

For control of brush and other woody plants, Imperial LO-VOL Super Brush Kill is the most effective herbicide available. It kills brush and other woody plants, and is safe for use on most crops and ornamentals. It is a low volatility herbicide, and it is safe for use on most crops and ornamentals. It is a low volatility herbicide, and it is safe for use on most crops and ornamentals.

DO NOT spray on crops or ornamentals. Do not spray on crops or ornamentals. Do not spray on crops or ornamentals.

DIRECTIONS

FOLIAGE (leaf) SPRAYING

Apply LO-VOL Super Brush Kill to woody growth in the afternoon. Do not spray until three weeks before fruiting. Best results are achieved from top to bottom. Apply in early summer when leaves are fully expanded. Do not spray in hot weather. Apply to the top and all parts of the plant. Do not spray stems and bark with spray. Any size brush can be treated. For best results, use a pump sprayer with nozzle. Do not use a backpack sprayer. Do not use a backpack sprayer.


DILUTION TABLE — LEAF SPRAYING

LO-VOL Super Brush Kill	Water
1 quart	100 gallons
1 pint	50 gallons
1 cup	25 gallons
1/2 cup	12 1/2 gallons
1/4 cup	6 1/4 gallons

Do not spray on crops or ornamentals. Do not spray on crops or ornamentals. Do not spray on crops or ornamentals.

BASAL TRUNK TREATMENT

Apply LO-VOL Super Brush Kill to the basal trunk of woody plants. Do not spray until three weeks before fruiting. Best results are achieved from top to bottom. Apply in early summer when leaves are fully expanded. Do not spray in hot weather. Apply to the top and all parts of the plant. Do not spray stems and bark with spray. Any size brush can be treated. For best results, use a pump sprayer with nozzle. Do not use a backpack sprayer. Do not use a backpack sprayer.



Imperial

TREAT:

- LEAVES
- STUMPS
- TRUNKS

LO-VOL SUPER BRUSH KILLER

CONTROLS:

- OAK
- HICKORY
- OSAGE ORANGE
- SUMAC
- and certain other woody plants
- ELM
- WILLOW
- PERSIMMON

A LOW VOLATILE HERBICIDE

ACTIVE INGREDIENTS	
**Mixed Octyl Ester of 2,4-Dichlorophenoxyacetic acid (minimum)	34.43%
**Mixed Octyl Ester of 2,4,5-Trichlorophenoxyacetic acid (minimum)	32.89%
INERT INGREDIENTS	32.68%
Total	100.00%

*Contains 2 pounds 2,4-Dichlorophenoxyacetic acid per gallon, equivalent to 22.85%
 **Contains 2 pounds 2,4,5-Trichlorophenoxyacetic acid per gallon, equivalent to 22.85%

CAUTION:
 KEEP OUT OF REACH OF CHILDREN
 SEE OTHER CAUTIONS ON SIDE PANEL

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

Manufactured by
IMPERIAL INC.
 SHENANDOAH, IOWA 51601

NET CONTENTS:



ST

Dilute LO-VOL Super Brush Kill for Basal Trunk Treatment. Wet the exposed roots. Do not spray on any time of day.

FR

Dilute LO-VOL Super Brush Kill for Basal Trunk Treatment. Wet the exposed roots. Do not spray on any time of day.

CAUTION: Avoid contact with eyes. Wash thoroughly if contact occurs.

DO NOT spray on crops and ornamentals.

Avoid spray on cotton, soybeans, clover, and food crops to drift. At the time of treatment, the same sprayer should be used for the same sprayer until the chemical is used. This product is not for use on crops and ornamentals. Do not use on crops and ornamentals.

APPLY ONLY to the basal trunk of woody plants. Do not spray on any time of day.

See the label for directions and precautions. Do not use on crops and ornamentals.

See the label for directions and precautions. Do not use on crops and ornamentals.

TREAT:

LEAVES
STUMPS
TRUNKS

Imperial

LO-VOL SUPER BRUSH KILLER

CONTROLS:

OAK
HICKORY
OSAGE ORANGE
SUMAG
ELM
WILLOW
PERSIMMON
and certain other woody plants

A LOW VOLATILE HERBICIDE

ACTIVE INGREDIENTS	
*Mixed Octyl Ester of 2,4-Dichlorophenoxyacetic acid (minimum)	34.43%
**Mixed Octyl Ester of 2,4,5-Trichlorophenoxyacetic acid (minimum)	32.89%
INERT INGREDIENTS	32.68%
Total	100.00%
*Contains 2 pounds 2,4-Dichlorophenoxyacetic acid per gallon, equivalent to 22.85%	
**Contains 2 pounds 2,4,5-Trichlorophenoxyacetic acid per gallon, equivalent to 22.85%	

CAUTION:

KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

Manufactured by

IMPERIAL INC.
SHENANDOAH, IOWA 51601

NET CONTENTS:

STUMP TREATMENT

Use Lo-Vol Super Brush Kill with the oil as the Base Treatment. Apply as coarse spray wet the tree, cut surface, bark and exposed roots down to the ground until the entire stump is saturated with the killing agent during the year.

FRILL TREATMENT

Use Lo-Vol Super Brush Kill with the oil as the Base Treatment. On trees over 6 inches in diameter in thick barked species cut overlapping notches in a ring around the trunk near base of trunk through bark but do not remove cambium. Apply to thoroughly saturate notches with the herbicide at any time of year.

CAUTION: HARMFUL IF SWALLOWED. Avoid contact with eyes, skin or clothing. In case of contact wash with soap and water.

DO NOT use around homes, recreational areas and similar sites.

Avoid spray drift to susceptible plants such as cotton, soybeans, tomatoes, grapes, sugarbeets, clover, alfalfa, ornamentals, fruit trees and food crops. Coarse sprays are less likely to drift. At high temperatures vapors from this product may injure susceptible plants in the immediate vicinity. Application by trunk treatment method during the dormant season will usually eliminate this hazard. Do not use the same spraying equipment for other purposes until thoroughly cleaned with a suitable herbicide cleaner. Do not store near fertilizers, seeds, insecticides or fungicides.

This product is toxic to fish. Avoid contamination of irrigation ditches, lakes, streams, ponds, etc. Do not use for domestic purposes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not mix with water by the use of a pump or sprayer.

APPLY ONLY AS DIRECTED ON THIS LABEL

Use only as directed on this label. Destroy it by burning or flushing it by or to a safe disposal away from water courses.

NOTICE

Read and understand the directions on the label and the brief directions on the back of any kit, container or label to which this label is attached and direct to the product for their use.

