DIRECTIONS FOR USE

VEGETABLE FOLIAGE INSECT CONTROL

Beans, Peas, Snap and Lima

Broccoli, Cauliflower, Cabbage

Carrots

Collards, Kale, Swiss Chard and Turnip Tops

Cucumbers

Lettuce, Endive, Aster, Aphids, Leaf Miners

Melons, Squash, Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelon, Persian and their hybrids, Watermelons and their hybrids, Summer and Winter Squash

Onions

Peas

Pepper

Imperial

DIAZINON® 12.5% E.C.

VEGETABLE SPRAY CONCENTRATE

ACTIVE INGREDIENTS:	
O,O-diethyl O-(2-isopropyl-6-methyl-4-primidinyl) phosphorothioate	11.5%
Aromatic petroleum derivative solvent	7.0%
INERT INGREDIENTS:	14.5%
Total	100.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER WARNINGS ON SIDE PANEL

WARNING—FLAMMABLE!
KEEP AWAY FROM HEAT OR OPEN FLAME

USE CONTENTS WITHIN 6 MONTHS OF OPENING — KEEP CLOSED

NET CONTENTS

Manufactured By
IMPERIAL INC.
SHENANDOAH, IOWA 51601

Radishes, Parsnips: Aphids, Flea beetle, Dipterous Leaf Miner. MIX 4 Tbs per 3 gallons water per 1,000 sq ft. Apply when insects appear and repeat at [illegible] Do not apply to radishes within 7 days of harvest.

Spinach, Beets: Aphids, Dipterous Leaf Miner. MIX 4 Tbs per 3 gallons water per 1,000 sq ft. Apply when insects appear and repeat at [illegible] Do not apply to spinach within [illegible] days of harvest. Do not apply to beets within 14 days of harvest.

Sweet Corn: Corn earworm. MIX 4 Tbs per 3 gallons water per [illegible] Apply when silk first appear and repeat applications [illegible] as necessary to 2-3 day intervals.

Tomatoes: Aphids, Dipterous Leaf Miner, Banded Cucumber Beetle, Fall Armyworm, Southern Armyworm. MIX 4 Tbs per 3 gallons water per 1,000 sq ft. Apply when insects first appear and repeat [illegible] Do not apply within 1 day of harvest.

VEGETABLE SOIL INSECT CONTROL

Cutworms, Mediterranean [illegible], Lettuce [illegible], Onion [illegible], Pea [illegible]

[illegible] fl oz per [illegible] sq ft [illegible]

Apply broadcast just before planting. Work into soil immediately 2-3 inches for surface cutworms, 6 inches for subsurface type.

Mole Crickets: [illegible] Cabbage, Corn, [illegible], Sweet Potatoes, Sugar Beets [illegible] Apply broadcast 1-2 days before planting and work into [illegible] inches after 1-2 days for mole crickets, 4-8 inches for wire worms.

CAUTION

Keep out of reach of children. Harmful if swallowed. May be absorbed through skin. Do not breath spray mist. Avoid contact with eyes and skin and clothing. Wash thoroughly after handling. Avoid contamination of food and feed.

Do not use on pets or humans. Do not allow children or pets on treated areas until spray has dried. Keep out of reach of domestic animals.

This product is toxic to fish and wildlife. Birds, fish or [illegible] may be killed. Keep out of lakes, ponds, or streams. Do not apply where runoff will drift from treated [illegible]. Do not apply where [illegible] Do not contaminate water by cleaning of equipment or disposal of wastes.

This product [illegible] toxic to bees exposed [illegible] Do not apply this product or allow it to drift to [illegible] Avoid the application [illegible] Apiarist or Pollinator Service.

Any use of this product in a manner not specified in this label [illegible]

NOTICE

[illegible] that the above [illegible] is safe or fit for the purpose [illegible] manufacturer [illegible] MERCHANTABILITY [illegible] A PARTICULAR PURPOSE [illegible] label [illegible]