DIRECTIONS FOR USE

VEGETABLE FOLIAGE INSECT CONTROL

Beans (Pole, Snap and Lima) ... Mix ... and Cucumber ... lbs 2 Tbsp ... water per 1,000 sq ft. Apply ... necessary. Do not apply within 7 days of harvest. Do not ... Do not apply within 4 days before grazing.

Broccoli, Cauliflower, Cabbage Aphids, Diamondback Moth, Imported Cabbageworm ... Mix 2 Tbsp per 3 gals. water per 1,000 sq ft. Apply ... Do not apply within 2 days of harvest. Do not ...

Carrots Aphids ... Mix ... Do not ...

Collards, Kale, Swiss Chard and Turnip Tops Aphids, Imported Cabbageworm, Diamondback Moth, Harlequin Cabbage Bug, Corn Earworm, Mix 2 Tbsp per 3 gals. water per 1,000 sq ft. Apply when ... Do not apply within 10 days of harvest. Do not apply ...

Cucumbers Aphids, Spotted Cucumber Beetle, Striped Cucumber Beetle, Mix ... Do not apply within 1 day of harvest.

Lettuce, Endive Aphids, Mix ... Do not apply within 7 days of harvest.

Melons (Squash) (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelon, Persian and their hybrids, Watermelons and their hybrids, Summer and Winter Squash) Aphids, Leafhoppers, Mites, Serpentine Leaf Miner, Thrips ... Mix 2 Tbsp per 3 gals. water per 1,000 sq ft. Apply ... Do not apply within 2 days of harvest.

Onions ... Thrip ... Mix ... per 1,000 sq ft. Apply ... Do not ...

Peas ... Mix ...

Peppers ... Mix ...

Potatoes ... Leaf Hopper ... Mix ...

Imperial

DIAZINON® 25% GARDEN CONCENTRATE

CONTROLS VEGETABLE GARDEN INSECT PESTS LISTED IN LABEL DIRECTIONS

ACTIVE INGREDIENTS:	
O,O-diethyl O-(2-isopropyl-6-methyl-4-primidinyl)phosphorothioate	25.0%
Aromatic petroleum derivative solvent	56.0%
INERT INGREDIENTS:	19.0%
Total	100.0%

WARNING

KEEP OUT OF REACH OF CHILDREN

SEE OTHER WARNINGS ON SIDE PANEL

WARNING — FLAMMABLE!
KEEP AWAY FROM HEAT OR OPEN FLAME

USE CONTENTS WITHIN 6 MONTHS OF OPENING — KEEP CLOSED

NET CONTENTS

Manufactured By

IMPERIAL INC.

SHENANDOAH, IOWA 51601

Radishes, Parsnips Aphids ... Mix ... per 1,000 sq ft. Apply when ... Do not apply within 10 days ...

Spinach, Beets Aphids, Downy Mildew, Leaf Miner ... Mix 2 Tbsp per 3 gals. water per 1,000 sq ft. Apply when ... Do not apply within 14 days of harvest. Do not apply ...

Sweet Corn Corn Earworm ... Mix 2 Tbsp per 3 gals. water per 1,000 sq ft. Apply when silk first appear ... Do not apply within 2 days of harvest.

Tomatoes Aphids, Downy Mildew, Leaf Miner, Broad Mites, Colorado Potato Beetle, Flea Beetle, Tomato Hornworm, Southern Armyworm ... Mix ... per 1,000 sq ft. Apply when ... Do not apply ...

VEGETABLE SOIL INSECT CONTROL

... surface and ... Price Beetle ... Wire Worms ... Mix ... Apply ... Do not ...

Mol ... Ants ...

WARNING

WARNING

NOTICE