

## DIRECTIONS FOR USE

### FIELD AND FORAGE INSECT CONTROL

Army Caterpillars, Spotted Armyworm, Yucca Flower Apterous, Pear Apterous, Leafhopper, Green Bug, Nymphs, Beetles, Aphids, and other insects which cause damage to crops by feeding on or sucking the sap from the stems, leaves, and fruit. Apply when a crop is first infested with these insects.

Grasshoppers. Apply 1 pint per acre to corn, sorghum, millet, and other crops. Do not apply to alfalfa or clover. Do not apply to alfalfa or clover if sheep, goats, or cattle are grazing immediately after application. Do not apply to alfalfa or clover if sheep, goats, or cattle are grazing immediately following cutting or mowing of hay provided direct or covered storage will not be made for 7 days following treatment.

Alfalfa Weevil Larvae. Apply 1 to 2 pints in sufficient water to give thorough coverage. Apply in spring when feeding damage to growing tips is evident. Repeat if necessary. If damage is extensive cut hay and apply to stubble. Do not graze dairy or beef cattle or sheep for 100 days. Do not cut for hay for 100 days. Do not treat in bloom. Remove stored hay during application.

Corn Field and Stem. Cut rootworm larvae. Planting time apply 2 1/2 to 3 1/2 oz. of seed in row of 47 inch long footed row in planter shoe in a minimum of 2 gallons water per acre. Post-emergence use: Treatment: Apply 1 1/2 to 2 1/2 fluid oz. per 1000 linear feet in sufficient water to wet base of corn plant and run a hand or soil next to plant. Corn treated area with 200 pounds of soil applied daily after spraying. Apply at first sign of injury.

Citronella. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

White Fly. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

Leaf Root Worm. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

Cotton Leaf Apterous. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

Grasshoppers. Apply 1 pint per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application. Do not apply to alfalfa or clover if sheep, goats, or cattle are grazing immediately following cutting or mowing of hay provided direct or covered storage will not be made for 7 days following treatment.

Cotton. Green Bug, Nymphs, Leafhopper, Legume Bug, Cotton Leaf Worm, Cotton Leaf Beetle, etc. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

Highway Caterpillars. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

Grasshoppers. Apply 1 pint per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

# Imperial

## DIAZINON® AG 4EC

### FOR THE CONTROL OF INSECT PESTS OF AGRICULTURAL CROPS

#### ACTIVE INGREDIENTS

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate **48.00%**

Xylene range solvent **38.96%**

INERT INGREDIENTS **13.04%**

Total **100.00%**

Contains 4 lbs. Diazinon per gallon  
Diazinon: Trademark Ciba Geigy Corporation

## WARNING

### KEEP OUT OF REACH OF CHILDREN

See Other Warnings on Side Panel

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

NET CONTENTS

Manufactured By

## IMPERIAL INC.

SHENANDOHE, IOWA 51601

## DIRECTIONS FOR USE

(Continued)

### FIELD AND FORAGE INSECT CONTROL

Grasshoppers. Apply 1 pint per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application.

Stink Bug, Midge. Apply 1 to 2 pints per acre to alfalfa. Do not apply to alfalfa if sheep, goats, or cattle are grazing immediately after application. Do not apply to alfalfa or clover if sheep, goats, or cattle are grazing immediately following cutting or mowing of hay provided direct or covered storage will not be made for 7 days following treatment.

Range land, pasture, grass forage, alfalfa, roadsides, noncrop lands, and barrier strip. Grasshoppers. Apply 1 to 2 pints per acre in sufficient water or oil for thorough coverage. When insects first appear, preferably in the nymph stage. Do not repeat for 30 days. Dairy or beef cattle or sheep may be grazed immediately following application, may be fed green forage immediately following cutting, and may be fed hay providing grass is not cut for 21 days following with or 30 days following spraying of animals.

#### WARNING

Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or clothing. Wash thoroughly after handling. Avoid contamination of food or feed. Keep out of reach of domestic animals. Do not use on lawns, lawns, or livestock. Keep out of reach of children or pets in treated areas until spray has dried.

This product is toxic to fish and aquatic life. Do not use in or near streams, rivers, ponds, lakes, or other bodies of water. Do not apply when treatment conditions could cause drift onto areas treated or untreated. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock.

Apply this product to lawns, lawns, or livestock. This product is toxic to fish and aquatic life. Do not use in or near streams, rivers, ponds, lakes, or other bodies of water. Do not apply when treatment conditions could cause drift onto areas treated or untreated. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock.

Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock.

#### NOTICE

This product is toxic to fish and aquatic life. Do not use in or near streams, rivers, ponds, lakes, or other bodies of water. Do not apply when treatment conditions could cause drift onto areas treated or untreated. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock. Do not use on lawns, lawns, or livestock.

Imperial Inc. is a registered trademark of Ciba-Geigy Corporation. © 1980 Ciba-Geigy Corporation.