

Imperial



2% DIAZINON[®] GRANULAR LAWN INSECT CONTROL

CONTROLS

Sod Webworm	Lawn Moths	Chiggers
Chinch Bugs	Millipedes	Crickets
Ants	Sowbugs	Digger Wasp
Brown Dog Ticks	Lawn Billbugs	Fleas
Clover Mites	Army Worms	Leafhoppers
Cutworms	Bermuda Mites	Springtails
Earwigs		Ticks

and

White Grubs of Japanese Beetles
European Chafers and Southern Chafers

12¹/₂ LBS. TREATS UP TO 2,500 SQUARE FEET

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 2⁰/₀

INERT INGREDIENTS 98⁰/₀

Total 100⁰/₀

Diazinon[®] Insecticide trademark of Ciba-Geigy

CONTROLS

Sod Webworm	Lawn Moths	Chiggers
Chinch Bugs	Millipedes	Crickets
Ants	Sowbugs	Digger Wasp
Brown Dog Ticks	Lawn Billbugs	Fleas
Clover Mites	Army Worms	Leafhoppers
Cutworms	Bermuda Mites	Springtails
Earwigs		Ticks

and

White Grubs of Japanese Beetles
European Chafer and Southern Chafer

12¹/₂ LBS. TREATS UP TO 2,500 SQUARE FEET

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 2⁰/₀

INERT INGREDIENTS 98⁰/₀

Total 100⁰/₀

Diazinon Insecticide trademark of Ciba-Geigy

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER CAUTIONS ON BACK PANEL

NET WEIGHT

Manufactured By

IMPERIAL INC.

SHENANDOAH, IOWA 51601

DIRECTIONS

OVERALL APPLICATION:

Diazinon Granules control the following insects on both grass and dichondra lawns: Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworm), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola).

Apply Diazinon Granules to your lawn with a fertilizer spreader at the rates shown below. Application should be made when grass is dry. Water lightly after application. Repeat as necessary.

SPREADER SETTING NO. (per 1,000 sq. ft.)

Spreader Make and Model	INSECTS LISTED BELOW*		White Grubs, Billbugs and Hyperodes Weevil 6 lbs
	Regular or Dichondra Lawn 5 lbs	St. Augustine Lawn 7 1/2 lbs	
Scott	6	7 1/2	6 3/4
Jackson 18-26 (18")	6	6 3/4	6 1/4
Sears Roebuck Cross Country (19")	5 3/4	6 3/4	6 1/4
Montgomery Ward 89-108 (20")	7 3/4	8 1/2	8 1/4
Salem Micro Set (22")	9	8 1/4	8 3/4
Cyclone Lawn Spreader	4 3/4	5 3/4	5 1/4
Lawn Beauty	6	6 3/4	6 1/4
Central Stamping	7 1/2	8	7 3/4

*Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworms), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola)

Note: Above spreader settings are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

For application with other types of equipment, or by means of a shaker, apply at the rate of 5 lbs. per 1,000 sq. ft. or 8 oz. per 100 sq. ft. For heavy lawns, application should be increased to 7 1/2 lbs. per 1,000 sq. ft. or 12 oz. per 100 sq. ft.

For: Billbugs and White Grubs of Japanese Beetle, European Chafer and Southern Chafer; Apply 6 lbs. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For Grub control make application any time between late July and early October. For Billbug control apply when activity is first noted or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

SPOT TREATMENT:

Ants: Apply 1 tablespoon of 2% Diazinon Granules over and around each hill.

Digger Wasps: Apply 1 tablespoon of 2% Diazinon Granules in and around each ground nest opening. Apply at dusk when insects are not active.

BAND TREATMENT AROUND FOUNDATIONS:

The following insects are commonly found around or near foundation of homes as well as in the lawn. One or more of these may even enter the home.

Ants	Clover Mites	Earwigs	Millipedes	Springtail
Brown Dog Ticks	Crickets	Fleas	Sowbugs	Collembola

Spreader Make and Model	INSECTS LISTED BELOW*		White Grubs, Billbugs and Hyperodes Weevil
	Regular or Dichondra Lawn 5 lbs	St Augustine Lawn 7 1/2 lbs	6 lbs.
Scott	6	7 1/2	6 3/4
Jackson 18-26 (18")	6	6 3/4	6 1/4
Sears Roebuck Cross Country (19")	5 3/4	6 3/4	6 1/4
Montgomery Ward 89-108 (20")	7 3/4	8 1/2	8 1/4
Salem Micro Set (22")	9	8 1/4	8 3/4
Cyclone Lawn Spreader	4 3/4	5 3/4	5 1/4
Lawn Beauty	6	6 3/4	6 1/4
Central Stamping	7 1/2	8	7 3/4

*Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworms), Bermuda Mites, Crickets, Army worms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola)

Note: Above spreader settings are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure correct rate of application.

For application with other types of equipment, or by means of a shaker, apply at the rate of 5 lbs. per 1,000 sq. ft. or 8 oz. per 100 sq. ft. For heavy lawns, application should be increased to 7 1/2 lbs. per 1,000 sq. ft. or 12 oz. per 100 sq. ft.

For: Billbugs and White Grubs of Japanese Beetle, European Chafer and Southern Chafer; Apply 6 lbs. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For Grub control make application any time between late July and early October. For Billbug control apply when activity is first noted or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

SPOT TREATMENT:

Ants: Apply 1 tablespoon of 2% Diazinon Granules over and around each hill.

Digger Wasps: Apply 1 tablespoon of 2% Diazinon Granules in and around each ground nest opening. Apply at dusk when insects are not active.

BAND TREATMENT AROUND FOUNDATIONS:

The following insects are commonly found around or near foundations of homes as well as in the lawns. Certain of these may even enter the home:

Ants	Clover Mites	Earwigs	Millipedes	Springtails
Brown Dog Ticks	Crickets	Fleas	Sowbugs	(Collembola)

For control apply a five foot band next to foundation. Distribute the granules uniformly using a rate of 8 oz. per each 5' by 20' area treated.

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contamination of food or feed. Avoid contact with skin, eyes or clothing. May be absorbed through skin. Wash thoroughly after using. Keep children and pets off treated area until material is washed into soil and grass is dry.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Apply only as specified on the label.

Do not reuse container. Destroy it by burying or burning. Avoid smoke fumes.