

Imperial*

5% DIAZINON[®] GRANULAR LAWN INSECT CONTROL

CONTROLS

Sod Webworm	Lawn Moths	Chiggers
Chinch Bugs	Millipedes	Crickets
Ants	Sowbugs	Digger Wasp
Brown Dog Ticks	Lawn Billbugs	Fleas
Clover Mites	Army Worms	Leafhoppers
Cutworms	Bermuda Mites	Springtails
Earwigs		Ticks

and

White Grubs of Japanese Beetles
European Chafer and Southern Chafer

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* 5%

INERT INGREDIENTS ----- 95%

Total ----- 100%

*Diazinon[®] Insecticide trademark of Ciba-Geigy

CONTROLS

Sod Webworm	Lawn Moths	Chiggers
Chinch Bugs	Millipedes	Crickets
Ants	Sowbugs	Digger Wasp
Brown Dog Ticks	Lawn Billbugs	Fleas
Clover Mites	Army Worms	Leafhoppers
Cutworms	Bermuda Mites	Springtails
Earwigs		Ticks

and

White Grubs of Japanese Beetles
European Chafer and Southern Chafer

ACTIVE INGREDIENT:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate* .. 5%

INERT INGREDIENTS 95%

Total 100%

* Diazinon® Insecticide trademark of Ciba-Geigy

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE OTHER CAUTIONS ON BACK PANEL

NET WEIGHT

Manufactured By

IMPERIAL INC.

SHENANDOAH, IOWA 51601

407-354

DIRECTIONS

OVERALL APPLICATION:

Diazinon Granules control the following insects on both grass and dichondra lawns: Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths (Sod Webworm), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola).

Apply Diazinon Granules to your lawn with a fertilizer spreader at the rate of 2-3 lbs. per 1,000 sq. ft. (an area 20' by 50') or 3.2-4.8 oz. per 100 sq. ft. (an area 10' by 10'). Application should be made when grass is dry. Water lightly after application. Repeat as necessary.

For application with other types of equipment, or by means of a shaker, apply at the rate of 2 lbs. per 1,000 sq. ft. or 3.2 oz. per 100 sq. ft. For heavy lawns, application should be increased to 3 lbs. per 1,000 sq. ft. or 4.8 oz. per 100 sq. ft.

For: Billbugs and White Grubs of Japanese Beetle, European Chafer and Southern Chafer; Apply 2 lbs. 6 oz. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

SPOT TREATMENT:

Ants: Apply 1½ teaspoons of 5% Diazinon Granules over and around each anthill.

Digger Wasps: Apply 1½ teaspoons of 5% Diazinon Granules in and around each ground nest opening. Apply at dusk when insects are not active.

BAND TREATMENT AROUND FOUNDATIONS:

The following insects are commonly found around or near foundations of homes as well as in the lawns. Certain of these may even enter the home:

Ants	Clover Mites	Earwigs	Millipedes	Springtails
Brown Dog Ticks	Crickets	Fleas	Sowbugs	(Collembola)

For control apply a five foot band next to foundation. Distribute the granules uniformly using a rate of 3.2 oz. per each 5' by 20' area treated.

CAUTION

sq. ft. (an area 20' by 50') or 3.2-4.8 oz. per 100 sq. ft. (an area 10' by 10'). Application should be made when grass is dry. Water lightly after application. Repeat as necessary.

For application with other types of equipment, or by means of a shaker, apply at the rate of 2 lbs. per 1,000 sq. ft. or 3.2 oz. per 100 sq. ft. For heavy lawns, application should be increased to 3 lbs. per 1,000 sq. ft. or 4.8 oz. per 100 sq. ft.

For: Billbugs and White Grubs of Japanese Beetle, European Chafer and Southern Chafer; Apply 2 lbs. 6 oz. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

SPOT TREATMENT:

Ants: Apply 1½ teaspoons of 5% Diazinon Granules over and around each anthill.

Digger Wasps: Apply 1½ teaspoons of 5% Diazinon Granules in and around each ground nest opening. Apply at dusk when insects are not active.

BAND TREATMENT AROUND FOUNDATIONS:

The following insects are commonly found around or near foundations of homes as well as in the lawns. Certain of these may even enter the home:

Ants	Clover Mites	Earwigs	Millipedes	Springtails
Brown Dog Ticks	Crickets	Fleas	Sowbugs	(Collembola)

For control apply a five foot band next to foundation. Distribute the granules uniformly using a rate of 3.2 oz. per each 5' by 20' area treated.

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contamination of food or feed. Avoid contact with skin, eyes or clothing. May be absorbed through skin. Wash thoroughly after using. Keep children and pets off treated area until material is washed into soil and grass is dry. Avoid breathing of dust.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Apply only as specified on the label.

Do not reuse container. Destroy it by burying or burning. Avoid smoke fumes.