

BAND TREATMENT AROUND FOUNDATIONS

The following insects are commonly found around or near foundations of homes as well as in the lawns. Certain of these may even enter the home:

Ants	Crickets	Millipedes
Brown Dog Ticks	Earwigs	Sowbugs
Clover Mites	Fleas	Springtails (Collembola)

For control apply a five foot band next to foundation. Distribute the granules uniformly using a rate of 3.2 oz. per each 5' by 20' area treated.

VEGETABLE INSECT PESTS

Vinegar flies (Drosophila spp.)

Tomatoes: Broadcast when flies first appear. Repeat in 7-10 day intervals or as required 2 oz. per 250 sq. ft. or $\frac{1}{2}$ to $\frac{3}{4}$ oz. per 100 linear ft., 10-20 lbs. per acre

Carrot Rust Flies

Carrots: Apply to seed furrow at planting. 4 oz. per 275 sq. ft. or $1\frac{1}{2}$ oz. per 100 linear ft., 40 lbs. per acre.

Cutworms (surface and subterranean)

Broccoli, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Cucumbers, Endive (escarole), Kale, Lettuce, Lima beans, Muskmelons, Parsley, Peas, Peppers, Potatoes, Snap beans, Spinach, Summer Squash, Sweet corn, Swiss chard, Tomatoes, Watermelons, Wintersquash: Apply broadcast just before planting. Work into soil immediately. 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. 4-8 oz. per 275 sq. ft. or $1\frac{1}{2}$ to 3 oz. per 100 linear ft., 40-80 lbs. per acre. For Potatoes mix into soil 4-6 inches deep.

Mole Crickets

Broccoli, Cabbage, Cauliflower, Collards, Endive (escarole), Kale, Lettuce, Mustard, Peppers, Radishes, Tomatoes, Turnips, Strawberries: Apply to plant beds or gardens 1-2 days before sowing seed or transplanting. Work into soil 1-2 inches immediately. 2 oz. per 275 sq. ft. or $\frac{1}{2}$ oz. per 100 linear ft., 20 lbs. per acre. For strawberries, do not apply within 5 days of harvest.

Onion Maggots

Onions: Apply in furrow with seed at planting. 2 oz. per 275 sq. ft. or $\frac{1}{2}$ oz. per 100 linear ft., 20 lbs. per acre.

Root Maggots

Broccoli, Brussels sprouts, Cabbage, Cauliflower: Apply broadcast and immediately work into soil 3-4 inches before seeding, either in seed bed or in garden or before transplanting. 4-6 oz. per 275 sq. ft. or $1\frac{1}{2}$ to 2 oz. per 100 linear ft., 40-60 lbs. per acre.

Radishes: Apply in seed furrow at planting. 3-4 oz. per 275 sq. ft. or $1\frac{1}{2}$ oz. per 100 linear ft., 20-40 lbs. per acre.

Turnips, Cabbage, Cauliflower, Broccoli: Apply in seed furrow at planting. 2 oz. per 275 sq. ft. or $\frac{1}{2}$ oz. per 100 linear ft., 20 lbs. per acre.

Wireworms

Broccoli, Cabbage, Cantaloupes, Carrots, Cauliflower, Celery, Collards, Cucumbers, Endive (escarole), Kale, Lettuce, Lima beans, Muskmelons, Onions, Parsley, Peas, Peppers, Snap beans, Spinach, Summer squash, Sweet corn, Sweet potatoes, Swiss chard, Tomatoes, Watermelons, Winter squash: Apply broadcast before planting and work into soil 4-8 inches immediately. 4-8 oz. per 275 sq. ft. or $1\frac{1}{2}$ -3 oz. per 100 linear ft., 40-80 lbs. per acre.

Potatoes: Apply at planting, placing in bands 3-6 inches deep along both sides of row. 2-4 oz. per 275 sq. ft. or $\frac{1}{2}$ to $1\frac{1}{2}$ oz. per 100 linear ft., 20-40 lbs. per acre

Corn (field and sweet)

Corn rootworm larvae (Diabrotica spp.): Apply in 7 inch bands centered over corn row at planting at the rate of 25-50 ounces per 1,000 linear feet of row.

Cutworm (surface and subterranean): For surface cutworms apply just before planting or when larvae first appear and work into soil 1-2 inches immediately. Do not apply later than layby.

For subterranean cutworms apply just before planting and work into soil 3-6 inches immediately. 4-6 oz. per 275 sq. ft. or $1\frac{1}{2}$ to 2 oz. per 100 linear ft., 40-60 lbs. per acre.

Fall armyworms: Apply over corn so granules fall into whorls when armyworms first appear. Repeat if necessary. 2-4 oz. per 275 sq. ft. or $\frac{1}{2}$ to $1\frac{1}{2}$ oz. per 100 linear ft., 20-40 lbs. per acre.

Southwestern corn borers: Apply over corn so granules fall into whorls when second generation borers first appear. Repeat one week later. 2-4 oz. per 275 sq. ft. or $\frac{1}{2}$ to $1\frac{1}{2}$ oz. per 100 linear ft., 20-40 lbs. per acre.

European corn borers: Apply so granules fall into whorls when 75% of plants show first generation borers feeding. Apply second application 7 days later if needed. For second generation borers, apply when field count shows 100 egg masses per 100 plants. 2-4 oz. per 275 sq. ft. or $\frac{1}{2}$ to $1\frac{1}{2}$ oz. per 100 linear ft., 20-40 lbs. per acre

Corn may be harvested immediately following last application. Do not feed treated forage to dairy or beef cattle or sheep within 10 days after foliar or over-the-corn applications exceeding $1\frac{1}{2}$ lbs. of active ingredient per acre

43

Imperial

5% DIAZINON GRANULAR INSECT CONTROL

CONTROLS

Ants	Cutworms	Millipedes
Army Worms	Digger Wasp	Mole Crickets
Bermuda Mites	Earwigs	Onion Maggots
Brown Dog Ticks	European Corn Borer	Root Maggots
Carrot Rust Flies	Fall Armyworms	Sow Bugs
Chiggers	Fleas	Springtails
Chinch Bugs	Lawn Billbugs	South Western
Cornroot Worm Larvae	Lawn Moths	Corn Borers
Clover Mites	(Sod Webworms)	Vinegar Flies
Crickets	Leafhoppers	Wireworms
	and	
	White Grubs of Japanese Beetles	
	European Chafer and Southern Chafer	

ACTIVE INGREDIENT:

O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate

INERT INGREDIENTS

Total

*Diazinon® Insecticide trademark Ciba-Geigy

KEEP OUT OF REACH OF CHILDREN CAUTION


STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN Wash immediately with soap and water.

IF IN EYES Flush eyes for at least 15 minutes with water. Call a physician immediately.



Manufactured By
IMPERIAL INC.
Shenandoah, Iowa 51601

NET WEIGHT

L-80-T-95778-K

44