

Phase 2-List C Response 1/4/90

Front Panel

IMPERIAL
PYRENONE 1-.2
FOOD PLANT SPRAY

A Ready-To-Use PYRENONE^R Insecticide

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF
MEAT AND POULTRY PROCESSING PLANTS

The insecticidal components piperonyl butoxide and pyrethrins are classified as food additives, and when this product is used as directed allowable tolerance levels will not be exceeded.

ACTIVE INGREDIENTS:

Pyrethrins	0.20%
*Piperonyl Butoxide, Technical	1.00%

INERT INGREDIENT:

Petroleum Distillate	<u>98.80%</u>
Total	100.00%

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% of related compounds.

PYRENONE^R is a registered trademark of Fairfield American

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 1 GALLON

EPA Reg. No. 407-355

E-74-84-1

EPA Est. 407-IA-1

Manufactured by
IMPERIAL INC.
SHENANDOAH, IA 51601

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticide Products and for Phase 2
Review, Page 2-12, As of 12/15/89 in Status.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food, or feed. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.

SPACE SPRAY: For rapid control of houseflies, fruit flies, mosquitoes, flying moths and gnats: Close doors and windows and shut off ventilating systems. Apply using a good sprayer adjusted to deliver a fine mist. Direct spray toward the upper portions of the enclosure, filling the room with mist. Use 1/2 to 1 ounce per 1,000 cubic feet of space.

To kill exposed stages of roaches, ants, spiders, clover mites, cheese mites, confused flour beetles, saw-toothed grain beetles, red flour beetles, rice weevils, granary weevils, and grain mites: Treat as above using 2 to 3 ounces per 1,000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY: To insure maximum effectiveness against the crawling insects named above, apply as a coarse mist directed toward insects, into cracks and crevices, (such as openings in expansion joints, between different elements of construction and between equipment and floors), along baseboards, around the base of machinery, containers of stored products and other areas where insects may hide or enter the building. Spray ant trails, nests and points of entry. Spray badly infested areas daily. In other areas, repeat treatment twice weekly or when reinfestation occurs.

NOTICE

Seller guarantee shall be limited to the terms of the label, and subject thereto, the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.