

Order the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 4407-35-9  
**ACCEPTED**  
 SEP 29 1975

# Imperial\*

# RABON® LIVESTOCK DUST


**A Ready-To-Use Insecticide Dust to Control  
 HORN FLIES AND LICE and aid in the control  
 of FACE FLIES on BEEF CATTLE  
 and LICE on SWINE**

<b>ACTIVE INGREDIENT</b>	<b>By Weight</b>
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	3.00%
<b>INERT INGREDIENTS</b> .....	<u>97.00%</u>
<b>Total</b> .....	<b>100.00%</b>

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
 See Other Cautions on Side Panel

NET CONTENTS \_\_\_\_\_

Manufactured By  
**IMPERIAL INC.**  
 SHENANDOAH, IOWA 51601



## GENERAL DIRECTIONS

RABON® Livestock Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

## USE DIRECTIONS

### BEEF CATTLE — HORN FLIES, FACE FLIES, LICE

**HAND DUSTING** — Apply approximately 2 oz. of dust by shake can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.

### SELF-TREATING DUST BAG

**FORCED USE** — Put RABON® Livestock Dust in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.

**FREE CHOICE USE** — Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice use aids in the control of lice.

### SWINE — LICE

**HAND DUSTING** — Apply 3-4 oz. of dust by conventional hand or powder duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

## PRECAUTIONS IN USING

**CAUTION:** Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings. Do not apply to animals less than one month old. Do not contaminate milk or milking equipment.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

## NOTICE OF WARRANTY

SELLER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

EPA Reg. No. 407-359  
EPA Est. 407-IA-1



O-4694B-H