

**NO. 2 + 2 LOW VOLATILE BRUSH AND WEED KILLER**

For control of mixed stands of brush consisting of sumac, elm, willow, poison ivy, and certain other softwoods as well as osage orange (hedge), hickory, oak, and other hardwoods. No. 2 + 2 Low Volatile Brush and Weed Killer is particularly effective for basal, stump, frill and foliage treatments in non-crop areas such as waste lands, fence rows, and for spot treatment in pastures.

DO NOT graze dairy animals on treated areas within six weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

**DIRECTIONS**

**FOLIAGE (leaf) SPRAYING**

Apply No. 2 + 2 Low Volatile Brush and Weed Killer to woody growth anytime after foliage is fully out until three weeks before frost. Best results are received from applications in late spring or early summer when leaves are fully out and before hot weather. Applications must be complete, and all parts of the plants, including foliage, stems and bark, wet with spray. Any size brush can be foliage sprayed, but basal or stump treatment will generally be more economical for brush over 8 to 10 feet high. For leaf spraying, mix only with water as shown below:


**DILUTION TABLE — LEAF SPRAYING**

No. 2 + 2 Low Volatile Brush and Weed Killer	Water
2 tablespoons	1 Gallon
2½ to 3½ cups	30 Gallons
2 to 3 pints	50 Gallons
2 to 3 quarts	100 Gallons

The lower dilution rates may be used when spraying just after the leaves are out. Repeat applications may be necessary if new growth develops, but a single treatment in any one season is usually sufficient. Broad-leaved weeds are also controlled by this application.

**BASAL (TRUNK) TREATMENT**

Dilute No. 2 + 2 Low Volatile Brush and Weed Killer at the rate of 1 pint to 3 gallons or 1 gallon to 25 gallons of fuel oil. Spray from the ground line up 18 inches on the trunk. Wet trunk on all sides and all exposed roots until run-off occurs. Use low pressure spray. Treat any time of year.



**NO. 2 + 2 LOW VOLATILE BRUSH AND WEED KILLER**

**CONTAINS BUTOXY PROPYL ESTERS OF 2,4-D AND 2,4,5-T**


**CONTROLS:**

**OAK HICKORY OSAGE ORANGE SUMAC and certain other woody plants**

**ELM WILLOW PERSIMMON**

**A LOW VOLATILE HERBICIDE**

<b>ACTIVE INGREDIENTS:</b>		
2,4-Dichlorophenoxyacetic Acid, Butoxy Propyl Esters		36.5%
2,4,5-Trichlorophenoxyacetic Acid, Butoxy Propyl Esters		34.5%
<b>INERT INGREDIENTS</b>		<b>29.2%</b>
<b>TOTAL</b>		<b>100.0%</b>
<b>CID EQUIVALENTS:</b>		
2,4-Dichlorophenoxyacetic Acid (2,4-D)	22.4%	— 2 Lb./Gal.
2,4,5-Trichlorophenoxyacetic Acid (2,4,5-T)	22.4%	— 2 Lb./Gal.
<b>4 POUNDS ACID EQUIVALENT PER GALLON</b>		



**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

See Other Cautions on Side Panel

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

**NET CONTENTS**

Manufactured By:

**IMPERIAL INC.**

SHENANDOAH, IOWA 51601

EPA Reg. No. 107-406  
EPA Est. 107-1A-1

**TREAT:**  
Leaves  
Stumps  
Trunks

**STUMP TREATMENT**

Dilute No. 2 + 2 Low Volatile Brush and Weed Killer with fuel oil as for Basal Treatment. Apply as coarse spray. Wet the freshly cut surface, bark, and exposed roots down to the ground until run-off occurs. Treat immediately after cutting any time during the year.

**FRILL TREATMENT**

Dilute No. 2 + 2 Low Volatile Brush and Weed Killer with fuel oil as for Basal Treatment. On trees over 6 inches in diameter or thick-barked species cut overlapping notches in a ring around the trunk near base. Cut through bark, but do not remove chips. Apply to thoroughly saturate freshly cut surfaces at any time of year.

**CAUTION: HARMFUL IF SWALLOWED.** Avoid contact with eyes, skin or clothing. In case of contact, wash with soap and water. DO NOT use around homes, recreational areas, and similar sites.

Avoid spray drift to susceptible plants such as cotton, soybeans, tomatoes, grapes, sugar beets, clover, alfalfa, ornamentals, fruit trees and food crops. Coarse sprays are less likely to drift. At high temperatures vapors from this product may injure susceptible plants in the immediate vicinity. Application by trunk treatment method during the dormant season will usually eliminate this hazard. Do not use the same spraying equipment for other purposes until thoroughly cleaned with a suitable chemical cleaner. Do not store near fertilizers, seeds, insecticides or fungicides.

This product is toxic to fish. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Avoid contamination of irrigation ditches, lakes, streams, ponds, ditch banks or water used for domestic purposes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. Coarse sprays are less likely to drift.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

**NOTICE**

Seller guarantee shall be limited to the terms of the label, and subject thereto, the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

N-316516-G