

[407-374]

DIRECTIONS FOR USE

Apply to dry foliage in the evening when air is calm. Begin application soon after plants are established or when insects or disease conditions threaten. Apply at 5 to 7 day intervals. Apply a light, even coating of dust to both upper and lower leaf surfaces.

IT IS IMPORTANT THAT FOLIAGE BE KEPT DUSTED, ESPECIALLY NEW GROWTH

TOMATOES: For anthracnose, early and late blights, grey leaf spot (stemphylium), septoria leaf spot — tomato fruitworm, yellow striped armyworm, aphids, spider mites, tomato russet mite, leafhoppers, apply at the rate of .1 lb. per 100 square feet (1.5 oz.). Do not exceed 40 lb. per acre. Do not use on tomatoes within 5 days of harvest.

CAUTION

Wash all vegetables thoroughly before eating. Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Avoid contamination of feed and food-stuffs.

ENVIRONMENTAL CAUTIONS

This product toxic to fish and wildlife.

Keep out of lakes, streams or ponds. Do not apply where run-off is likely to occur.

Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply only as specified on this label.

This product is toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

Do not reuse container. Destroy when empty.

NOTICE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Imperial

TOMATO GARD

Insecticide and Fungicide for Tomatoes

ACTIVE INGREDIENTS:

*Captan	5.0%
**Malathion	4.0%
INERT INGREDIENTS:	91.0%
Total	100.0%

*N-[(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide

**O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Other Cautions on Back Panel

EPA Reg. No. 407-374
EPA Est. 407-IA-1

O-1210B-D

<p>ACCEPTED</p> <p>MAY 9 1975</p> <p>Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, this pesticide registered under EPA Reg. No. 407-374</p>	<p>CONTENTS _____</p> <p>Manufactured by</p> <p>IMPERIAL INC.</p> <p>SHENANDOAH, IOWA 51601</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------