

407-374

PM 14

1062

Front Panel

IMPERIAL

TOMATO GARD

Insecticide and Fungicide for Tomatoes

ACTIVE INGREDIENTS:

.....	5%
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	4%
INERT INGREDIENTS:	91%
Total	100%

~~.....~~ Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 407-374
EPA Est. 407-IA-1


E-75-84-1

NET CONTENTS

Manufactured by
IMPERIAL INC.
SHENANDOAH, IA 51601

ACCEPTED
 JUL 31 1986
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 407-374

BEST AVAILABLE COPY



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Wash thoroughly after using. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Avoid contamination of feed and foodstuffs. Wash all vegetables thoroughly before eating.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply only as directed on this label.

This product is toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not allow persons to enter treated areas within 4 days following application unless protective clothing is worn. Conspicuously post reentry information at site of application.

Do not rotate a foliar treated crop with crops other than those with registered captan uses.

Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

STORAGE AND DISPOSAL

STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food or feed. Keep container closed.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

Apply to dry foliage in the evening when air is calm. Begin application soon after plants are established or when insects or disease conditions threaten. Apply at 5 to 7 day intervals. Apply a light, even coating of dust to both upper and lower leaf surfaces.

IT IS IMPORTANT THAT FOLIAGE BE KEPT DUSTED, ESPECIALLY NEW GROWTH.

TOMATOES: For anthracnose, early and late blights, grey leaf spot (stemphylium), septoria leaf spot, and the insects tomato fruitworm, yellow striped armyworm, aphids, spider mites, tomato russet mite, leafhoppers, apply at the rate of .1 lb. per 100 sq ft of harvest.

NOTICE: Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herein.

BEST AVAILABLE COPY