

Front Panel

TOMATO GARD

Insecticide and Fungicide for Tomatoes

ACTIVE INGREDIENTS:

| | |
|---|---------------|
| *Captan | 4.9% |
| Related Derivatives | 0.1% |
| Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate | 4.0% |
| INERT INGREDIENTS: | <u>91.0%</u> |
| Total | 100.0% |

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 407-374
EPA Est. 407-IA-1

I-86-84-1

NET CONTENTS _____

Manufactured by
IMPERIAL INC.
Shenandoah, IA 51601

per self cert pfm
 NOV 12 1987
 407-374

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Wash thoroughly after using. Do not get in eyes. Avoid contact with skin and clothing. Avoid contamination of feed and foodstuffs. Wash all vegetables thoroughly before eating.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician. If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply only as directed on this label.

This product is toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.

Wear chemical resistant gloves when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing.

Do not contaminate water by cleaning of equipment or disposal of wastes.
STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food, or feed. Keep container closed.
PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

Apply to dry foliage in the evening when air is calm. Begin application soon after plants are established or when insects or disease conditions threaten. Apply at 5 to 7 day intervals. Apply a light, even coating of dust to both upper and lower leaf surfaces.

IT IS IMPORTANT THAT FOLIAGE BE KEPT DUSTED, ESPECIALLY NEW GROWTH.

TOMATOES: For anthracnose, early and late blights, grey leaf spot (stemphylium), septoria leaf spot, and the insects tomato fruitworm, yellow striped armyworm, aphids, spider mites, tomato russet mite, leafhoppers, apply at the rate of .1 lb. per 100 square feet (1.5 oz.). Do not use on tomatoes within 7 days of harvest.

NOTICE: Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.