

Imperial

RABON[®] LIVESTOCK, POULTRY & PREMISE SPRAY

50% WETTABLE POWDER INSECTICIDE

ACTIVE INGREDIENTS:	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphate	50.0%
INERT INGREDIENTS	50.0%
Total	100.0%



CAUTION

KEEP OUT OF REACH OF CHILDREN

See other cautions on side panel.

GENERAL DIRECTIONS

RABON 50 WP is suitable for use in conventional power or low pressure knapsack sprayers. Occasional agitation is recommended to prevent undue settling of the suspension. Follow the USE DIRECTIONS for the proper percent solution needed for a specific insect and areas or types of wall surfaces. Refer to DILUTION TABLE for the quantity of insecticide needed to make the percent solution recommended.

Do not apply to animals less than one month old.

PRECAUTIONS IN USING

CAUTION: Harmful if swallowed. Avoid breathing spray mist and dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees. Avoid direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NET CONTENTS

Manufactured by
IMPERIAL INC.
SHENANDOAH, IOWA 51601

LIVESTOCK USE DIRECTIONS

ANIMAL	INSECT	% SOLUTION	REMARKS
Beef Cattle	House Flies	1.0%	Apply to walls, ceiling, floor cracks, and crevices using a power sprayer.
	Stable Flies	1.0%	Apply to walls, ceiling, floor cracks, and crevices using a power sprayer.
Swine	House Flies	1.0%	Apply to walls, ceiling, floor cracks, and crevices using a power sprayer.
	Stable Flies	1.0%	Apply to walls, ceiling, floor cracks, and crevices using a power sprayer.

POULTRY USE DIRECTIONS

INSECT	TYPE OF HOUSING	% SOLUTION	REMARKS
Flies & Mites	Wire Cages	0.5%	Apply evenly to birds of Gal. 1000 Part. 1.0% solution at 1.0% off areas from below. Repeat when necessary. Do not repeat more often than every 10 days. For individual birds, immerse in solution of the mixture for 10 days.
	Floor Management Litter	0.5%	Apply 1/2 Gal. 1000 Sq. Ft. evenly for penetration to litter surfaces. Also apply thoroughly to walls, floor cracks, crevices and interior surfaces for 10 days.
	Floor Management Litter	5.0% WP	Treat evenly to 1.0% thoroughly using 1/2 Oz. 1000 Sq. Ft.
	Floor Management Dust Box	5.0% WP	Mix evenly throughout top layer of box contents using 2 1/2 Oz. 1000 Birds.
	Floor Management Roost Paint	1.0%	Treat with brush or sprayer thoroughly to walls, ceiling, floor cracks, and crevices using 1/2 Pt. 1000 Ft.
Powl Tick (Blue Bug)	All Types	1.0%	Apply 1 Gal. 1000 Sq. Ft. thoroughly to walls, ceiling, floor cracks, and crevices with a power sprayer.

Note: For maximum long control of the northern fowl mite, penetration of the feathers around the vent area is absolutely vital. Use a power sprayer at 100-125 p.s.i. at NO LESS THAN RECOMMENDED RATE. More attention must be given each individual bird when using low pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks.

PREMISE USE DIRECTIONS

INSECT	PROBLEM AREAS	% SOLUTION	REMARKS
Flies	Dairy barns, poultry houses, swine barns - other animal buildings.	2.0%	For dry, whitewash, wood or concrete, use 1/2 Gal. 1000 of 2.0% solution.
		1.0%	For painted wood or other smooth black surfaces, use 1/2 Gal. 1000 of 1.0% solution.
		1.0%	For masonry or galvanized, heat metal surfaces, use 1/2 Gal. 1000 of 1.0% solution.
Maggots	Poultry droppings, manure piles, garbage piles, animal feed troughs.	1.0%	Apply 1/2 Gal. of 1.0% solution to 100 sq. ft. of manure area for 10 days. Repeat every 10 days thereafter.
Ticks	Campgrounds, backyards, pastures, etc. (residential areas, etc. - outdoor areas only).	1.0%	Spray 1/2 Gal. 1000 thoroughly on yard, pastures, etc. of 1.0% solution. Repeat every 10 days. For pastures, repeat every 10 days.

DILUTION TABLE

AMOUNT OF RABON [®] WETTABLE POWDER	AMOUNT OF WATER FOR			
	0.35% SOLUTION	0.5% SOLUTION	1.0% SOLUTION	2.0% SOLUTION
1/2 lb.	9 Gal.	6 Gal.	3 Gal.	1 1/2 Gal.
2 lb.	36 Gal.	24 Gal.	12 Gal.	6 Gal.
4 lb.	72 Gal.	48 Gal.	24 Gal.	12 Gal.
8 lb.	144 Gal.	96 Gal.	48 Gal.	24 Gal.

NOTICE OF WARRANTY

SELLER MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

EPA Reg. No. 105-1001
EPA Est. No. 107-1A-1

© 1974 A