DIRECTIONS FOR USE

Do Not Exceed Application Rate of 50 lb. Per Acre

TOMATOES: Foliage Application - To control Anthracnose, early and late blight, grey mold (cladosporium), grey leafspot (stemphy'ium), nailhead spot, septoria leafspot. sou hern blight (sclerotium) and the insect pest; Colorado potato beetle, flea beetle, leafhoppers, climbing cutworms. European corn borer, fall armyworm, lacebugs, pinworms, stinkbugs, tomato hornworm, tomato fruitworm, tarnished plant bug. To control Colorado potato beetle, flea beetles, leafhonpers and climbing cutworms, apply at the rate of % oz. per 100 square feet; 9 oz. per 1000 square feet; or 25 lb. per acre. For other listed pests, apply at the rate of 1 % oz. per 100 square feet; 1 lb. 2 oz. per 1000 square feet; 50 lb. per acre. For disease control, begin when first fruit clusters form or disease threatens. Apply at 7-10 day intervals more frequently under severe disease conditions. Do not apply within 5 days of harvest.

PEPPERS: To control anthracnose, bacterial leafspot, downy mildew, cercospora and other leafspots, phytophthora and other fruit rots. and the insects Colorado potato beetle, flea beetles, leafhoppers, climbing cutworms, corn earworm. European corn borer, fall armyworms, stinkbugs, tomato hornworm, tarnished plant bug. To control Colorado potato beetle, flea beetle, leafhoppers and climbing cutworms, apply at the rate of % oz. per 100 square feet; 9 oz. per 1000 square feet; or 25 lb. per acre. To control other listed pests. apply at the rate of 1% oz. per 100 square feet; 1 lb. 2 oz. per 1000 square feet; or 50 lb. per acre. For disease control, begin when fruit first forms or disease threatens. Apply at 7-10 day intervals or more frequently under severe disease conditions.

EGGPLANT: For the control of alternaria leafspot (early blight), begin in plant bed; anthracnose, cercospora leafspot, downy mildew, Phomopsis and other fruit rots, phytophthora (late blight) and the insects Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lacebugs, pinworms, stinkbugs, tomato fruitworms, tomato hornworms, tarnished plant bugs. For Colorado potato beetle, flea beetle, and leafhoppers, apply at the rate of % oz. per 100 square feet; 9 oz. per 1000 square feet; or 25 lb. per lore. For other listed pests, apply at the rate of 1% oz. per 100 square feet; 1 lb. 2 or 1000 square feet; or 50 lb. per acredisease control, apply when fruits first or disease threatens. Repeat at 7-10 day increased under tervals — more frequently under sever sease conditions.

CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Wash hands and face before eating and smoking. Take shower or bath after using. Wear long-sleeved clothing. Do not store near or contaminate feed or foodstuffs.

ENVIRONMENTAL CAUTIONS


This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of waste. Do not app. where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not use except as specified on this label.

Do not reuse container. Destroy by burning or burying. Stay out of smoke or fumes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

NOTICE

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with direct us given herein.



For Disease and Insect Control

ACTIVE INGREDIENTS: Zineb (zinc ethylene bisdithiocarbamate) 6.0 % * Carbaryl; 1-naphthyl methylcarbamate 4.0 % INERT INGREDIENTS 90.0 % Total 100.0 %

*Metallic Zinc equivalent 1.42%

CAUTION

KEEP OUT OF REACH OF CHILDREN
See Other Precautions on Side Panel

NET WEIGHT

EPA Reg. No. 407-388 EPA Est. 407-IA-1 O-4167B-C

Manufactured by

IMPERIAL INC.

SHENANDOAH, IOWA 51601