

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION
Harmful if swallowed. Avoid inhalation of spray mist. Wash thoroughly after handling. Wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other approved State and local procedures.

CROP	PRODUCT	WATER
FRUITS		
apples, apricots, cherries, nectarines, peaches, pears, plums, prunes and quinces	4 oz. ----- 1 pint -----	1 gal. 4 gal.
BERRIES		
blackberries, boysenberries, dewberries, grapes, loganberries, raspberries, blueberries, cranberries, currants, gooseberries, and strawberries	4 oz. ----- 1 pint ----- 3 gal. 2 oz. -----	1 gal. 1 gal. 100 gal.
NUTS		
almonds, filberts, pecans and walnuts	4 oz. ----- 1 pint ----- 3 gal. 2 oz. -----	1 gal. 4 gal. 100 gal.

Always dilute with water. The spray tank must be free of any sulfur residues before using this product. Do not combine sulfur sprays with this product or follow applications of sulfur for at least 14 days. Do not use as a summer spray or if green foliage is showing. Do not spray evergreens, hickory, or maple trees with this product. Retrate frequently while spraying.

Imperial

DORMANT SPRAY OIL
For Mite, Scale and Leafroller Control

ACTIVE INGREDIENTS:
Petroleum Oil 90%
NET INGREDIENTS 10%
Total 100%

*Petroleum oils minimum unsulfonated residue 90%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swalloweds: Do not induce vomiting. If vomiting occurs keep airway open. Seek medical aid.
If on Skin: Wash with soap and water.
If in Eyes: Flush with water for 15 minutes. Seek medical aid.

See Other Precautions on Side Panel

EPA Reg. No. 407-392 Manufactured by: with COMMENTS-G
EPA Est. 407-1A-1 **IMPERIAL INC.** EPA Letter Date:

SHERMANSBURG, IOWA 51601 407-392

NET CONTENTS DEC 1 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this product is registered under EPA Reg. No.

APPLES: To control European Lecanium scale, oystershell scale, To control European red mite, scale, San Jose scale, cottony European red mite, Forbes scale, shell scale, San Jose scale, white peach scale, fruittree, Jose scale, and olive scale. To control European red mite, Forbes scale, scurfy scale, To control European red mite, oystershell scale, San European red mite, Forbes scale, cottony peach scale or mite, Forbes scale, fruittree Jose scale, cottony peach scale, red mite, Forbes scale, oyster ply of the rate of 1/2 cup D per 4 gallons of water. Cover of peach leaf curl on peaches as prepared above at the rate of

BLACKBERRIES, BOYSENBERRIES and RASPBERRIES: To control two-spotted spider mite. BL spotted spider mite. CRAMB To control two-spotted spider control two-spotted spider mite and two-spotted spider of water; or 1 pint per 4 gallons of water.

ALMONDS: To control olive red spider mite, fruittree le European red spider mite, P and obscure scale. WALNUT European red spider mite. Apply at the rate of 1/2 cup of water; or 3 gallons 2 ou dormant spray before growth s weather when spray can dr

Seller guarantee shall be l thereto, the buyer assumes use or handling and accept

BEST DOCUMENT AVAILABLE



**STATEMENTS
DOMESTIC ANIMALS**

pray mist. Wash thoroughly after use.

HAZARDS
Do not apply directly to lakes, streams, or rivers. Do not apply when needed. Do not contaminate water courses. Apply this product only as

DR USE
product in a manner inconsistent

DISPOSAL
by storage or disposal. Open

or rinse water that cannot be disposed of according to approved methods. Then offer for recycling or by landfill or by other approved

PRODUCT	WATER
4 oz.	1 gal.
1 pint	4 gal.
4 oz.	1 gal.
1 pint	4 gal.
3 gal. 2 oz.	100 gal.
4 oz.	1 gal.
1 pint	4 gal.
3 gal. 2 oz.	100 gal.

must be free of any sulfur residues sulfur sprays with this product or 4 days. Do not use as a summer spray evergreens, hickory or quantity while spraying.

Imperial

DORMANT SPRAY OIL For Mite, Scale and Leafroller Control

ACTIVE INGREDIENTS:	
Petroleum Oils*	99%
INERT INGREDIENTS	1%
Total	100%

*Petroleum oils minimum unsulfonated residue 90%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Do not induce vomiting. If vomiting occurs keep always open. Seek medical aid.
Eyes: Flush with water for 15 minutes. Seek medical aid.
Skin: Wash with soap and water.

See Other Precautions on Side Panel

EPA Reg. No. 407-392
EPA Est. 407-1A-1

ACCEPTED
IMPERIAL INC.
SHEMANGOAN, IOWA 51001

NET CONTENTS
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended by the Federal Insecticide, Fungicide, and Rodenticide Act, FIFRA Reg. No. 407-392

APPLES: To control European red mite, Forbes scale, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale, scurfy scale. **APRICOTS:** To control European red mite, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale, cotony peach scale, olive scale. **CHERRIES:** To control European red mite, Forbes scale, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale and brown soft scale. **NECTARINES:** To control white peach scale, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale, and olive scale. **PEACHES:** To control white peach scale, European red mite, Forbes scale, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale, scurfy scale, cotony peach scale, and olive scale. **PEANUTS:** To control European red mite, Forbes scale, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale, scurfy scale. **PLUMS:** To control European red mite, Forbes scale, lecanium scale, oystershell scale, San Jose scale, cotony peach scale and olive scale. **PRAIRIE:** To control European red mite, Forbes scale, fruitree leafroller, lecanium scale, oystershell scale, San Jose scale, cotony peach scale and olive scale. **RASPBERRIES:** To control European red mite, Forbes scale, oystershell scale, San Jose scale and scurfy scale. Apply at the rate of 1/2 cup Dormant Spray Oil per gallon of water or 1 pint per 4 gallons of water. Cover entire tree thoroughly. For supplement control of peach leaf curl on peaches, add Ferbam 76% WP to the spray solution as prepared above at the rate indicated on the Ferbam label.

BLACKBERRIES, BLYTHEBERRIES, BOWENBERRIES, GRAPES, LOGANBERRIES, and RASPBERRIES: To control European red mite, Pacific spider mite and two-spotted spider mite. **BLISSBERRIES:** To control lecanium scale and two-spotted spider mite. **CRANBERRIES:** To control lecanium scale. **CURRENTS:** To control two-spotted spider mite and San Jose scale. **GOOSEBERRIES:** To control two-spotted spider mite. **STRAWBERRIES:** To control Pacific spider mite and two-spotted spider mite. Apply at the rate of 1/2 cup per gallon of water; or 1 pint per 4 gallons of water; or 3 gallons 2 ounces per 100 gallons of water.

NUT CROPS

ALMONDS: To control olive scale, oystershell scale, San Jose scale, European red spider mite, fruitree leafroller and spider mite. **PILBERTS:** To control European red spider mite. **PECANS:** To control San Jose scale, spider mite, and obscure scale. **WALNUTS:** To control oystershell scale, San Jose scale, European red spider mite, spider mite, lecanium scale and walnut scale. Apply at the rate of 1/2 cup per 1 gallon of water; 1 pint per 4 gallons of water; or 3 gallons 2 ounces per 100 gallons. Apply as a delayed dormant spray before growth starts in the spring and during a period of mild weather when spray can dry before freezing. Cover entire tree thoroughly.

NOTICE

Seller guarantee shall be limited to the terms of the label, and subject thereto, the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

BEST DOCUMENT AVAILABLE

DEC 1 1982

Mr. G. E. Currie
Imperial, Inc.
P.O. Box 423
Shenandoah, IA 51001

Dear Mr. Currie:

Subject: ~~Imperial Product 2000-11~~
~~2000-11~~
Deletion of Label Text
Application dated November 4, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (78-767)

Enclosure

RD:ELLENBERGER:DCR-28969:WANG-1314C:pjb:Raven:479-2013:11/24/82