

HAZARDS TO HUMANS
DANGER

ENVIRONMENTAL HAZARDS

STORAGE AND DISPOSAL

TRAFFIC
REGULATION

DIRECTIONS FOR USE

MIXING TABLE

APPLICATOR
EQUIPMENT

Table of Equivalents

ACCEPTED
Imperial
DEC 16 1980

Approved for the Federal Insecticide, Fungicide, and Plant Growth Regulator Act

LINE SULPHUR SPRAY

INSECTICIDE AND FUNGICIDE FOR FRUIT TREES
HARDY TREES, SHRUBS, ROSES AND BERRIES

ACTIVE INGREDIENTS	10.00%
Carrier, Propellant	70.00%
INERT INGREDIENTS	100.00%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

STATEMENT OF PRACTICAL TREATMENT
1. Mix with water in the ratio of 1 part of spray to 100 parts of water.
2. Apply to foliage of trees and shrubs for 15 minutes.
3. Get more information from the manufacturer.
4. See the label for additional cautions.

PACKED BY
IMPERIAL INC.
CHENANDOSH, IOWA 51601

NET CONTENT

DCRMANT SPRAY

APPLICATOR EQUIPMENT AND CHERRIES

FRUIT

FRUIT TREES AND SHRUBS

**EARLY SPRING SPRAYING
AS GROWTH STARTS**

FRUIT

FRUIT TREES AND BLACKBERRIES

SUMMER FOLIAGE SPRAY

FRUIT

FRUIT

FRUIT TREES

ACCEPTED
DEC 16 1980
Imperial

Under the Federal Insecticide, Fungicide, and Fungicide Act, as amended, and the Federal Food, Drug, and Cosmetic Act, as amended.

LIME SULPHUR SPRAY

INSECTICIDE AND FUNGICIDE FOR FRUIT TREES
SHADE TREES SHRUBS ROSES AND BERRIES

ACTIVE INGREDIENT	30.0%
Inert Ingredients	70.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

1. On Skin: Wash with soap and water.
2. In Eye: Flush with plenty of water for 15 minutes.
3. Get Medical Attention.

SEE THE PANEL FOR ADDITIONAL CAUTION

PACKED BY

IMPERIAL INC.

CHENANDUAK, IOWA 51601

NET CONTENTS

DORMANT SPRAY

EARLY SPRING SPRAYING
AS GROWTH STARTS

SUMMER FOLIAGE SPRAYING