


SPACE SPRAYS: fruitflies, gnats, grain moths, house flies, hornets, mosquitoes, skipper flies, wasps. Dilute one part in 59 parts water (2 tablespoons in 3 1/4 pints water). Use at the rate of 1/2 ounce diluted spray per 100 cubic feet of space. Direct the space treatment upward. Keep windows and doors closed during treatment and for at least 10 minutes after application.

VAPORIZER APPLICATORS: To kill fruitflies, gnats, Angoumois grain moths, house flies, hornets, mosquitoes, skipper flies and wasps where oil residues are not undesirable, dilute with deodorized base oil instead of water and apply at the dilution rate of 1 part to 59 parts by means of any good vaporizer type applicator.

NOTICE

Seller guarantee shall be limited to the terms of the label, and subject thereto, the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.



4

4107-4107

PM 17

1/12/90

123

Front Panel

**IMPERIAL
PYRENONE®
MULTI-PURPOSE SPRAY**

**A Concentrated PYRENONE®
Micro Emulsion**

**Clear - Stable - Leaves No Oily or
Objectionable Residues**

Effective, Fast Acting Insecticide

**For Use in Homes, Institutions, Restaurants
Food Processing Plants, Dairies, Livestock and
Outdoors for Mosquito Control**

**Apply with Conventional Mechanical or
Compressed Air Equipment**

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	1.0%
INERT INGREDIENTS:	98.5%
Petroleum Distillate	0.4%
Total	100.0%

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

PYRENONE® - Registered Trademark of Fairfield American Corp.

KEEP OUT OF REACH OF CHILDREN

CAUTION


SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 1 QUART

**EPA Reg. No. 407-409
EPA Est. 407-IA-1**

K-82-84-1

**Manufactured by
IMPERIAL INC.
SHENANDOAH, IA 51601**



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After space spraying in food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food, or feed. Keep container closed. Do not transport or store under 32°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). The offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatments. All food processing surfaces should be covered during treatment or thoroughly cleaned before use.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drug-store beetles, meal worms, grain mites and caddisflies: Use a good sprayer, adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers or stored foods, around the base of machinery, behind shelves and drawers. For silverfish spray bookcases. For ants spray trails, nests, and points of entry. If surface application only is to be used spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make

