

• localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

**To control fleas and brown dog ticks:** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices along and behind baseboards, mouldings, window and door frames and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

**To control bedbugs:** Spray mattresses lightly particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

**To control clothes moths:** Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediteranean flour moths, Indian meal moths and tobacco moths: Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3 - 6 ounces per 1,000 cubic feet of space.

**FOGGING:** To kill exposed or accessible stages of the above named crawling insects use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

**ON ANIMALS:** To protect cattle and horses from horn flies, house flies, mosquitoes and gnats apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies apply at a rate of 2 ounces per adult cow or horse sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.


**OUTDOOR APPLICATION**

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools or ponds and shore lines. Apply at a rate of 1 gallon per acre (4 ounces per 1,000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist models 51C91 or 100HT traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 45 gallons per minute.


**NOTICE**

Buyer assumes all risks of use, storage or handling of this product in strict accordance with directions given herewith.



Front Panel

IMPERIAL  
PYRENONE R LOW OIL  
DAIRY SPRAY



A Concentrated PYRENONE R Micro Emulsion

Effective, Fast Acting Insecticide

Clear - Stable

Leaves No Oily or Objectionable Residues

Apply with Conventional Mechanical  
or Compressed Air Equipment

ACTIVE INGREDIENTS:

Pyrethrins .....	0.1%
*Piperonyl Butoxide, Technical .....	1.0%
INERT INGREDIENTS: .....	98.5%
Petroleum Distillate .....	0.4%
Total .....	100.0%

\*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

PYRENONE R Registered Trademark of Fairfield American Corp.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 407-410  
EPA Est. 407-IA-1

K-82-84-1

NET CONTENTS  
Manufactured by  
IMPERIAL INC.  
SHENANDOAH, IA 51601

Not for use on...  
...  
...