

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Remove pets, birds, and cover fish aquaria before spraying.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a secure area, in original container only, away from fertilizers, food, or feed. Do not transport or store under 32°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

ON ANIMALS: To protect cattle and horses from horn flies, mosquitoes, and gnats apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies apply at a rate of 2 ounces per adult cow or horse sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3 - 6 ounces per 1,000 cubic feet of space.

NOTICE

Buyer assumes all risks of use, storage, or handling of this product not in strict accordance with directions given herewith.

Phase 2-List B Response 1/5/90

Front Panel

IMPERIAL
PYRENONE^R DUST

Control Cockroaches, Crickets, Fruit Flies,
And Many Common Garden Pests.

Leaves No Objectionable Residues

For Use in Residences, Institutions, Food Plants, Canneries,
On Stored Seed And Grain
And On Vegetable And Ornamental Plants.

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
*Piperonyl Butoxide, Technical	1.00%
INERT INGREDIENTS:	97.00%
Petroleum Distillate	1.90%
Total	100.00%

*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl)
ether and 0.2% related compounds.

PYRENONE^R Registered Trademark of Fairfield American Corp.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS


NET CONTENTS _

EPA Reg. No. 407-411
EPA Est. 407-IA-1

L-82-84-1

Manufactured by
IMPERIAL INC.
SHENANDOAH, IA 51601

[Faint, illegible text]



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing dust. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a secure area, in original container only, away from fertilizers, food, or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

FOR CONTROL OF COCKROACHES, CRICKETS
SILVERFISH, FIREBRATS AND CENTIPEDES

Using conventional dusting equipment, apply this product at 2 ounces per 100 square feet to all indoor areas where these pests may crawl or hide. Apply in cracks and crevices and hidden surfaces around, under and behind baseboards, cabinets, refrigerators, stoves, sinks, doors, pipes, windows, electrical fittings, electrical equipment and other possible harborages. Repeat application as required.

ON VEGETABLE AND ORNAMENTAL PLANTS

Pyrenone Dust may be used throughout the growing season; or as a pre-harvest treatment. Acceptable for use up to the day of harvest on edible plants when other insecticides are prohibited due to residue restrictions and use limitations. Apply dust uniformly to upper and lower leaf surfaces at the rate of 30 pounds of dust per acre. Treat only when the insects are present and repeat as required to maintain control.

ORNAMENTALS - To control aphids, flea beetles, leafhoppers, white flies on African Violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, dogwoods, geraniums, gladiolus, marigolds, roses, etc. **ASPARAGUS** - To control asparagus beetles. **BEANS** - To control aphids, leafhoppers, white flies, Mexican bean beetles. **BROCCOLI, CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER** - To control aphids, cabbage loopers, cross-striped cabbageworms, diamondback moth larvae, flea beetles, harlequin bugs, imported cabbageworms. **CELERY** - To control green peach aphids, leaf hoppers, cabbage loopers. **CRANBERRIES** - To control fireworms, leafhoppers. **EGGPLANT** - To control Colorado potato beetles, flea beetles, green peach aphids. **LETTUCE** - To control cabbage loopers, green peach aphids, diamondback moth larvae, imported cabbageworms. **PEPPERS** - To control green peach aphids. **POTATOES** - To control aphids, Colorado potato beetles, flea beetles, leafhoppers. **RADISH** - To control aphids and flea beetles. **SPINACH** - To control aphids, cabbage loopers, and webworms. **TOMATOES** - To control green peach aphids, flea beetles, Colorado potato beetles.