

July 16, 1987  
Front Panel

IMPERIAL

READY-TO-USE  
RAT & MOUSE KILLER

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

11 MAR 1988

Contains Warfarin -- Cheese Flavored  
For Control of Rats and Mice

ACTIVE INGREDIENT:

Warfarin (3-alpha-Acetylbenzyl)-4-  
hydroxycoumarin .....

0.025%

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

INERT INGREDIENTS: .....

99.975%

407-435

Total ..... 100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 407-UGL

EPA Est. 407-IA-1

Manufactured by  
IMPERIAL INC.  
Shenandoah, IA 51601

NET CONTENTS \_\_\_\_\_

Side/Back Panel

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

## ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container, in a cool, dry area inaccessible to children and pets. Keep container closed.

PRODUCT DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

USE RESTRICTIONS: For control of Norway rats, roof rats, and house mice in and around homes, industrial and agricultural buildings and similar man-made structures and inside transport vehicles (ships, barges, railcars, aircraft).

SELECTION OF TREATMENT AREAS: Determine areas where rats/mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much of any other food source as possible.

## APPLICATION DIRECTIONS:

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals, or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

FOR RATS -- Place 4 to 16 oz. of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

FOR MICE -- Place 1/4 to 1/2 oz. (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

*replace* Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is near, such as a dump or uncontrolled area such as an abandoned building, establish permanent bait stations and replenish bait as needed.

NOTICE: Buyer assumes all risk of use, storage, or handling of this product not in strict accordance with directions given herewith.