

C3793

SUGGESTIONS FOR USE

BEEF CATTLE: For control of Horn Flies, Lice, and Ticks: Mix 1 pint with 19 gallons water (1 gallon with 150 gallons water). Apply as a spray to wet animal thoroughly. One spraying should destroy adult lice and some eggs, providing long lasting protection; kill Horn Flies present and provide several weeks residual kill; kill adult ticks quickly and provide residual protection for several weeks. May be repeated at three week intervals as needed.

SHEEP AND GOATS: For control of Lice, Keds (Sheep Ticks), fleeceworms, and Ticks: Mix 1 pint with 19 gallons water (1 gallon with 150 gallons water). Apply as a spray for lice, Keds, and ticks. A single thorough wetting will usually be sufficient for practical control: For Fleeceworms, wet infested and surrounding area thoroughly. Repeat as necessary.

HOGS: For control of lice: Thoroughly mix 1 pint with 20 gallons water. Spray animals thoroughly. Apply only one time. Do not spray animal within 30 days of slaughter. Do not use on pigs less than 3 months old. Do not treat sows within two weeks of farrowing nor for at least 3 weeks thereafter.

BACKRUBBER INSECTICIDE: For control of Horn Flies and as an aid in control of lice and face flies on beef cattle. Mix 1 pint with 9 pints diesel fuel (1 gallon with 9 gallons). Apply to backrubber until thoroughly saturated. When used in commercial type backrubbers or applicators, care should be taken to see that oil is not dispensed too freely as excessive quantities may "scald" animals in extremely hot weather. For lice control, backrubber should be of a type that will permit application of insecticide to all parts of the body. Prompt control of Horn Flies may be expected, but several weeks of continuous use will be necessary to lower louse population. For face fly control, animals must regularly rub face and head. Backrubber should be constructed so that it is readily available for rubbing of the face and head. Animals under six months of age should not be exposed to the back rubber treatment.

Backrubber insecticides are more effective in hot weather when prepared with a mixture made up of 1 part 30W motor oil (non-detergent) and 2 parts diesel fuel. This heavier solution is less volatile keeping the insecticide more effective for a longer time. In hot weather backrubbers should be treated often for the most effective results.

TOXA

An Err

HC
LICE, KED

To control H

Active Ingredients:
*Technical Chlo
Gamma Isomer
Petroleum Der
(Aromatic
Inert Ingredients ...

NET CON

WARN

FRAN

STICKER

SPREADER

Sept 16 1971

410 + 4

41730

41717

C3793



TOXAPHENE—LINDANE Spray Mix

An Emulsifiable Insecticide For Use In Making:

A SPRAY TO CONTROL

HORN FLIES, LICE, AND TICKS on BEEF CATTLE

LICE, KEDS (Sheep Ticks) AND TICKS on SHEEP AND GOATS

A BACK-RUBBER INSECTICIDE

To control HORN FLIES and aid in controlling FACE FLIES and LICE on
BEEF CATTLE

Active Ingredients: Toxaphene*	45%
*Technical Chlorinated Camphene (Chlorine content 67-69%)	
Gamma Isomer of Benzene Hexachloride from Lindane	2%
Petroleum Derivative Solvents	38%
(Aromatic, 15%; Aliphatic, 23%)	
Inert Ingredients	15%

NET CONTENT: 128 FL. OZ. (ONE GALLON)

WARNING: KEEP OUT OF REACH OF CHILDREN

See Other Warnings of Side Panel
USDA REGISTRATION NO. 410-44

WARNING

May be fatal if swallowed. Vapor and spray mist may be absorbed through the skin. Keep out of eyes. Do not contact with skin or clothing. In the event contact is made, wash thoroughly. Do not breathe fumes or spray mist.

Do not use or store in or around the home

Do not use Toxaphene-Lindane on dairy animals, in dairy barns or milk rooms. Do not use on meat animals within 30 days of slaughter. Do not use on extremely young animals (under 3 months). Do not exceed concentrations suggested on label. Do not use on any animals not included in specific recommendations on this label. Avoid spraying livestock during excessively cold or hot weather. Do not spray overheated or visibly sick animals, and avoid working or long drives following spraying. This product is not recommended for use as a dip.

Do not permit contamination of feed or foodstuffs.

Do not use in poultry houses.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands, taking care not to contaminate water supplies. Containers should be disposed of by breaking or by punching holes in them and burying with wastes, or by burning. Keep out of smoke.

This product is poisonous to wildlife. Birds feeding on treated areas may be killed.

Do not apply where runoff is likely to occur. Keep out of rivers, streams, and ponds.

Apply this product only as specified on this label.

POISON

Call a Physician at Once! ANTIDOTE: If swallowed, give a tablespoonful of salt (sodium chloride) in a glass of warm water and repeat until vomit is clear. Wash thoroughly with soap and water to remove from skin or clothing.

Prepared by
FRANKLIN LABORATORIES INC.
DENVER, COLORADO 80222, U.S.A.