


C3351



LIVESTOCK INSECTICIDE POWDER

**An Insecticidal Dusting Powder
for Control of**

CONTAINING:

Active Ingredients:

*Methoxychlor, Technical	5%
Melathion	4%
(O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	

Inert Ingredients	91%
-------------------	-----

*5% Methoxychlor, Technical is equivalent to 4.4% 2, 2-bis (p-Methoxyphenyl)-1, 1, 1-trichloroethane and 0.6% other isomers and reaction products.

CAUTION

Do not use on very young animals (under 6 weeks of age).

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. After application, wash hands with soap and water.

Do not contaminate feed, milk, water or utensils.

Toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse container. Destroy when empty.

NET WEIGHT: 16 Oz. (1 Pound)

See side panel for additional cautions

EPA Registration No. 410-48-AA

Horn Flies and Face Flies: For hand application, apply one to two heaping tablespoonsful of Livestock Insecticide Powder to the animal, taking care to cover poll, neck, entire topside and upper parts of the side. Rub powder lightly into the hair, to the skin, but do not rub into the skin. For face flies, apply powder to animal's face and rub lightly into hair. Repeat as needed.

Horn Flies and Face Flies: Apply as described above for beef cattle.

CAUTION: Do not apply to lactating dairy animals within 5 hours of milking or during milking.

Repeat hand application to lactating dairy animals no more frequently than every three weeks.

Apply as above to entire body, with particular attention to those areas where parasites congregate. For control of hog lice, powder may be scattered throughout bedding, being careful not to contaminate feedstuffs or water.

Franklin Livestock Insecticide Powder is available in 5 pound packages and 50 pound drums

EPA Est. 410 (C) 1