

DIRECTIONS FOR USE

Screwworm and Blowfly Infestations: Apply with swab or brush, working into deep pockets. Also spread smear around and over drainage area below wound. Re-treat if necessary. Freely bleeding wounds may require several applications.

To Prevent Screwworm and Blowfly Infestations: Treat fresh wounds resulting from dehorning, castrations, shear cuts, etc. Also treat navels of newborn animals. Re-treat as needed.

Fleece Worm and Other Blowfly Infestations: Mix one part Blue Smear Screwworm Killer with nine parts water. Apply as a wash on and around infested areas.

Ear tick and ear mite control: Pour uniluted smear in ears or apply with a brush or swab.

EPA EST. 410-CO-1

C3703



**BLUE SMEAR
SCREWWORM KILLER**

CONTAINS KORLAN*

Kills screwworms, ear ticks, ear mites and blowfly maggots in cattle, sheep, swine and horses of all ages, including newborn animals.

ACTIVE INGREDIENTS:

Ronnel — O,O-dimethyl O-(2,4,5-trichlorophenyl) phosphorothioate (from technical grade)..... 5%
Xylene 5%

INERT INGREDIENTS 90%

EPA Registration No. 410-74-AA

For External Use Only

LIVESTOCK REMEDY — NOT FOR HUMAN USE

NET CONTENT: 14 FL. OZ.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

See Side Panel for Additional Directions

CAUTION

Avoid breathing of vapor. Do not leave in eyes or on human skin, wash thoroughly. Avoid contamination of feed and foodstuffs. May be absorbed through the skin. Harmful if swallowed.

Do not reuse container. Destroy when empty.

*Trademark of The Dow Chemical Company.

Prepared by
FRANKLIN LABORATORIES INC.
DENVER, COLORADO 80222, U.S.A.

71214