

C3421

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or clothing. Wash thoroughly with soap and water after handling and before eating or milking. Avoid contamination of feed and foodstuffs. Do not apply in dwellings.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

GENERAL DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of in a sanitary landfill, or by incineration if allowed by state or local authorities.

Franklin Louse-Fly Killer is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

POULTRY USE DIRECTIONS

For birds housed in wire cages, apply 1 lb./300 birds by plunger, rotary type duster, or shaker can duster. For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than once every 14 days. Wire rungs and corners should also be treated.

NOTE: For Northern Fowl Mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.



Louse-Fly Killer

A Dusting Powder for Control of HORN FLIES and LICE on Beef and Dairy Cattle and Horses, LICE on Swine, STABLE FLIES, HORSE FLIES, GNATS, DEER FLIES, HOUSE FLIES, and TICKS on Horses, NORTHERN FOWL MITES, and LICE on or around Chickens, and as an aid in the control of FACE FLIES on Beef and Dairy Cattle and Horses.

Active Ingredient:
2-chloro-1-(2,4,5-trichlorophenyl)
vinyl dimethyl phosphate* 3.00%
Inert Ingredients 97.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

If Swallowed — Call a physician immediately.
If In Eyes — Wash with plenty of water for 15 minutes; if irritation persists, see a physician.
If On Skin — Wash thoroughly with soap and water.
See Side Panel For Additional Precautionary Statements.


*Rabon® — Registered Trademark of Shell Chemical Company
EPA Registration No. 410-80 EPA Est. No. 410-CO-1

Net Wt. 1 lb. (453 g)

83381C



Franklin Laboratories, Inc.
Aurora, Colorado 80044



POULTRY FLOOR MANAGEMENT

LITTER: Apply 1 lb./100 sq. ft. by plunger or rotary type duster. Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.

DUST BOX: Use 2 lb./100 birds, in a box about 2 ft. by 2 ft. x 1 ft. deep.

ROOST PAINT: Use 1 lb./100 ft. — Make thick slurry by mixing 1 lb. of dust with 1 pint of water, stirring continually. Treat roosts thoroughly, particularly the cracks and crevices.

LIVESTOCK USE DIRECTIONS

BEEF AND DAIRY CATTLE: Horn Flies, Face Flies and Lice.

HORSES: Horn Flies, Stable Flies, Horse Flies, Lice, Gnats, Deer Flies, Face Flies, House Flies, Ticks.

HAND APPLICATION: Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Wearing a dust mitt, rub in lightly to carry the dust beneath the hair. Repeat as necessary.

SWINE: Lice
DUSTER APPLICATION: Apply 3-4 oz. of dust by conventional and/or powder duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

DUST BAG APPLICATION

FORCED USE: Put Franklin Louse-Fly Killer in cotton cloth or double burlap bags or use pre-packed weatherproof cattle dust bags and hang in the barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.

FREE CHOICE USE: Use in the same dust bags as above, but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice use aids in the control of lice.

TAKE TIME

