

AGRICULTURAL CHEMICAL
 Do not mix with Food, Feeds, Drugs, or Clothing
 Do not use on animals with open wounds, cuts, or sores
 Do not use on animals with open wounds, cuts, or sores
 Do not use on animals with open wounds, cuts, or sores
 Do not use on animals with open wounds, cuts, or sores



DURSBAN 44 INSECTICIDE

FOR CONTROL OF LICE AND HORN FLIES ON
 BEEF-BREED CATTLE ONLY

Active Ingredients:
 Chlorpyrifos (90-Diethyl U) (5.6-Inchlore-2) 43.2%
 pyridyl phosphorothioate) 56.8%
 Total Ingredients: 100%
 Contains 3.8 pounds of Chlorpyrifos per gallon
 Contains petroleum distillates

EPA Reg. No. 410-82 EPA Est. No. 77 NY 1

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS
 MAY BE FATAL IF SWALLOWED • MAY BE ABSORBED THROUGH SKIN
 • MAY BE IRRITANT TO EYES AND SKIN
 • Do Not: Take Internally • Do Not: Use in Eyes or on Skin
 • Wear: No Car Gloves When Handling • Wash Thoroughly After Handling
 • Wash Contaminated Clothing Before Reuse • Avoid breathing vapors.

STATEMENT OF PRACTICAL TREATMENT
 If Swallowed • Do not induce vomiting • Call a physician immediately if on skin • In case of contact, remove contaminated clothing and immediately wash skin with soap and water. If in Eyes • Flush eyes with plenty of water and get prompt medical attention.
 Note to Physician and Veterinarian • Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS
 This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS
 COMBUSTIBLE • Do not use near open flame or heat.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PRECAUTIONS: Do not contaminate water, food or feed by storage or disposal. Open flame is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide or waste that cannot be used or chemically repackaged should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternate procedures such as incineration by burning.

FRANKLIN DURSBAN 44 Insecticide is a convenient "one shot" treatment to provide season-long control of biting and irritating FRANKLIN DURSBAN 44 is not effective against cattle gnats. Cattle may be treated during any season of the year for lice and horn fly control. FRANKLIN DURSBAN 44 provides for control of horn flies up to 6 weeks.

HOW TO APPLY: Wearing rubber gloves, and in a well ventilated area, pour the required dose of FRANKLIN DURSBAN 44 in one spot to the top middle area of the animal, just behind the shoulder blades and neck junction.

AMOUNT TO USE: Use FRANKLIN DURSBAN 44 for lice and horn fly control at the rate of 2cc per 100 lb of animal body weight up to 800 lb. For animals weighing more than 800 lb, use no more than 16cc. Due to the small amount of material required, care must be taken to apply the proper dose.

Animal Weight	Amount of DURSBAN 44 Administered/Animal	No. of Animals Contents of this Container will Treat
Do Not Treat Animals Under 200 Lbs.		
200 lb	4cc	118
300 lb	6cc	78
400 lb	8cc	58
500 lb	10cc	47
600 lb	12cc	39
700 lb	14cc	33
800 lb	16cc	29
Over 800 lb	16cc	29

USE PRECAUTIONS
 DO NOT TREAT • Bulls over 8 months of age of any breed • Purbred continental or exotic breed cattle such as Simmental, Charolais, Charolais or Gelshorn • Dairy breed cattle of any age • Cows within 21 days prior to or 14 days after calving • Veal calves • Beef breed calves under 12 weeks of age
 Dairy, Brauns, continental or exotic breed cattle crossed with beef breeds (Brauns) may be treated.
 Do not treat sick, convalescent or severely stressed animals.
 Do not treat cattle for 10 days before or after shipping, detaching, castration, vaccination, etc.

Do not slaughter animals within 14 days of when treatment is reapplication does occur, etc. It may be re-treated according to the following schedule: (1) 45 days after initial treatment with a 14 day withdrawal prior to slaughter or (2) 30 days after initial treatment with a 21 day withdrawal prior to slaughter. Do not re-treat animals within less than 30 days.

FRANKLIN DURSBAN 44 contains a cholinesterase inhibitor. Do not use any drug or other peridical chemical having cholinesterase inhibiting activity, either simultaneously or within 45 days before or after treatment with it.
 The solvent system in FRANKLIN DURSBAN 44 evaporates quite rapidly which may cause a cold sensation to the animal at the site of application. This may manifest itself as a very mild transitory irritation to the animal.
 Do not brand animals with a hot iron while treating with FRANKLIN DURSBAN 44. The solvent system is combustible and could cause a serious burn to the animal.
 US Patent No. 3244576

NOTICE: Seller warrants that the product contains no chemical deterioration and is reasonably fit for the purposes stated. The user when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Manufactured by
FRANKLIN LABORATORIES, INC.
 Denver, Colorado 80222

*Trademark of THE DOW CHEMICAL COMPANY.

ACCEPTED

MAR 27 1981

U.S. Dept. of Agriculture
 Plant Industry and Research Administration
 Agricultural Research Service
 Registered under
 EPA Reg. No. 410-82