



PIONEER

DIRECTIONS

ARMYWORMS—On Wheat, Oats, Barley, Sorghum and Rye, use $2\frac{2}{3}$ pints in 5 to 10 gallons of water per acre. See Caution Statement.

CUTWORMS—Variiegated, Black, and Subterranean Cutworms on Corn, Sorghum, use $2\frac{2}{3}$ pints in 10 gallons water per acre. Spray to give uniform application. For those working below ground, direct nozzles over row, apply in late evening. Cultivate dirt over row after treatment. See Caution Statement.

GRASSHOPPERS—In Corn, Sorghum, Wheat, Barley, Rye and Fence Rows, use 2 to $2\frac{2}{3}$ pints in 10 gallons of water per acre. Use low rate for immature grasshoppers. Spray to give uniform coverage. See Caution Statement.

NON WARRANTY

Bartels and Shores warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purpose. Since timing, methods of application, weather, plant, and soil conditions, mixture with other chemicals and other factors affecting the use of this prodct are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

EMULSIFIAB
TOXAPHEN
CONCENTRA
Effective Control of Army
Cutworm and Grasshop

BARTELS



WARNING STATEMENT:

Do not feed treated forage, hay or corn ensilage to dairy animals or animals being finished for slaughter.

Do not apply to forage to be sold commercially or shipped interstate.

No limitation on use of harvested grain.

Do not ensilage treated forage.

Do not apply within 2 weeks of harvest on barley; 1 week on wheat, oats and rye; 4 weeks on grain sorghum.

Do not apply to sorghum more than once after heads start to form.

Do not graze dairy animals being finished for slaughter in treated fields.

WARNING: May be fatal if swallowed! Do not breathe spray mist. Do not get on skin, clothing or in eyes. Wash thoroughly after using. Do not contaminate feed or foodstuffs.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.

Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place.

Not for use or storage in or around the home.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.



PIOR

EMUL
TOXA
 CON
 Effective Co
 Cutworm

CONTENTS:
1 QUART

Contains 6 l
ACTIVE INGREDIENTS:
 Toxaphene (Techn
 containing 67
 Insecticidally activ
INERT INGREDIENTS:

EPA REG. NO. 413-64

FEDERAL INSECT
 ND RODENTICID
 IC POISON REGI

Handwritten signature

1400 ST. LOUIS A