


[416-54]



[416-54]



1 lb. **QUINN'S** Net

ONE PER CENT ROTENONE GARDEN DUST

FOR THE CONTROL OF CERTAIN
INSECTS ON VEGETABLES

ACTIVE INGREDIENTS:

Rotenone _____ 1.00%
Other cube' resins _____ 2.00%

INERT INGREDIENTS: _____ 97.00%

Read entire label. Use in strict accordance with label directions and cautions.

Caution: Keep out of reach of Children

Avoid inhalation of dust. Do not get in eyes. Wash or wipe edible portions of vegetables thoroughly before use.

Manufactured By
Quinn Drug & Chemical Company
GREENWOOD, MISSISSIPPI

ENVIRONMENTAL CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

EPA Reg. No. 416-54 EPA Est. 416-Ms-1

DIRECTIONS FOR USE:

AS A DUST:

Dust thoroughly at rate of one pound per 1000 square feet (30-40 pounds per acre). Start application as soon as insects appear and repeat at 5-10 day intervals unless otherwise specified, or as necessary, and in accordance with recommendations of local county agents, experiment stations and extension service. Do not apply within 1 day of harvest. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

AS A SPRAY

Mix 6-8 tablespoonfuls in a little water to make a thin paste. Add paste to one gallon water and agitate during application.

BEANS—Mexican bean beetle, thrips, flea beetle.

CABBAGE, COLLARDS, TURNIPS, and MUSTARD—Cabbage worms, flea beetle.

CUCUMBER, SQUASH, and CANTALOUPE—Cucumber beetle, flea beetle, melon worms

PEAS—Pea aphids

POTATOES—Colorado potato beetle, potato flea beetle

TOMATOES—Tomato horn worm when small, tomato suck fly, flea beetle