

Cenol
PROLIN.

Cenol

ACCEPTED

RATS AND MICE

ORDER
FUNGIC
FOR E
BY UN
AT

WARFARIN RODENTICIDE

**KILLS
RATS-MICE**

Cenol
PROLIN
BAIT
STATION

1. GENERAL DIRECTIONS

- a. Place baits in locations where rats and mice feed, water, or travel.
- b. Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.
- c. Contaminated or spoiled baits should be replaced promptly.
- d. Baiting should continue until all signs of feeding have stopped.

2. DIRECTIONS FOR RAT CONTROL

- a. Four ounces of bait per station should be the minimum amount used.
- b. Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.
- c. An uninterrupted supply of bait should be maintained for at least 15 days.

3. DIRECTIONS FOR MOUSE CONTROL

- a. Tablespoon amounts (1/2 to 1 ounce) of bait should be placed at 8-12 foot intervals.
- b. Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.
- c. An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION

Protect children, pets, and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this product may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K, combined with blood transfusions, are indicated, as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

Baits should be placed in areas inaccessible to children, pets, and domestic animals, or in tamper-proof bait boxes.

To use as bait station
cut carton on perforated
lines to permit
easy feeding.

ACTIVE INGREDIENTS:

Warfarin	0.025%
2-(4-chlorophenyl)-5-hydroxy-1,4-dioxane	0.025%
N-(4-chlorophenyl)-2,6-dimethyl-4-pyridinamine	0.025%
Calciferrolol	0.025%

INERT INGREDIENTS

99.950%
100.000%

SEE BACK
FOR DIRECTIONS

FORMULATED AND PACKED BY
CENOL COMPANY, INC.
CHICAGO 51, ILLINOIS

Cenol
PROLIN

Cenol PROLIN... with bait

Cenol PROLIN
Ready To Use
MICE STATION

Cenol

Cenol



ACTIVE INGREDIENTS

0.025

0.025

INERT INGREDIENTS

99.950

100.000

Satisfaction
Guaranteed

Satisfaction
Guaranteed

CAUTION - KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTION ON BACK PANEL

Cenol

... with bait
Ready To Use

12-3-71
419-142

FOR USE IN SUBJECT
ED UPON COMMENTS.

Pull down perforated triangles on the sides of the box. Place bait in locations where mice feed, water or travel. Place Prolin Mice Stations along walls, in corners, and in concealed locations where mice can feed unobserved are desirable. Contaminated or spoiled baits should be replaced promptly.

PLACE CENOL PROLIN MICE STATIONS AT 8-12 FOOT INTERVALS.

BAITS SHOULD BE PLACED IN AREAS INACCESSIBLE TO CHILDREN, PETS, WILDLIFE AND DOMESTIC ANIMALS OR IN TAMPER PROOF BAIT BOXES.

An uninterrupted supply of bait should be maintained for at least 15 days.

CAUTION

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

Baits should be placed in areas inaccessible to children, pets and domestic animals or in tamper proof bait boxes.

U.S.D.A. REG. #419-142