FRONT PANEL

CENOL
WARFARIN
CONCENTRATE

ACTIVE INGREDIENT:

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER PRECAUTIONS ON BACK PANEL.

Weight 4 - 8 oz.

ACCEPTED

APR 17 1968

UNDER THE TIME AND MET CICIDE FUNGILITE AND ME ONT CIDE ACT FOR FORMOVIC FOLION REGISTERATED UNDER NO. 17-13 BULLICT TO ATTACHED COMMENTS.

Manufactured By

Cenol Company, Chicago, Illinois

DIRECTIONS

RATS

Mix thoroughly one part of Warfarin Concentrate with 19 parts of a suitable bait material which is available, such as corn meal, rolled oats, dog food, laying mash, stock food, etc. A small amount of a bland oil, such as mineral, corn, peanut, or any other edible oil, may be used, up to 5%, when the bait is so dry that it might have a tendency to dust.

When perishable baits, such as ground meat, fish, vegetables or similar materials are used, bait placements should be made in late afternoon and replaced daily with fresh bait. Dogs and cats will actively search out baits of this type. Therefore, baits must be placed so as to prevent feeding of these animals.

Baits should be placed in amounts of from 4 oz. up to 1 lb. in locations frequented by rats or where rats have been known to feed. Baits must be protected from children, dogs, cats, poultry, livestock, etc., by bait boxes, cages or other suitable means. An uninterrupted bait supply should be maintained for at least 10 days.

Baits should be examined regularly and replaced when completely eaten. If bait moulds, decomposes or becomes sour it should be replaced.

MICE

For controlling mice follow the same general directions as for rats except that the amount of bait placements should be smaller (1 tablespoonful) and more placements should be made at 8 - 12 foot intervals.

Mice are more difficult to control than rats and an uninterrupted supply of baits should be maintained for at least 15 days.