

Cenol

SEVIN 50

WETTABLE POWDER

INSECTICIDE

ACTIVE INGREDIENT:

\*Carbaryl (1-naphthyl N-methylcarbamate)

50.0%

INERT INGREDIENTS:

50.0%

100.0%

\*Sevin, U.S. Pat. Nos. 2,903,478; 3,009,855

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Harmful if inhaled or swallowed. Avoid prolonged breathing of dust or spray. Do not take internally. Skin contact may be harmful. Avoid prolonged or repeated contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following spraying.

Do not use with lime, Bordeaux, lime sulfur, casein-lime spreaders or other alkaline materials.

Application of Cenol Sevin 50W to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

Apply this product only as specified on this label.

NET WEIGHT: 12 CZ.

Registered under EPA Reg. No. 419-162

Cenol Sevin 50 Wettable Powder is to be dispersed in water using constant agitation in the spray tank during application. This product may be applied within one day of harvest on all food crops listed unless otherwise indicated. Observe intervals between last application and harvest indicated by number of days in ( ) following crop.

FOR USE EAST OF THE ROCKY MOUNTAINS FOR APPLES AND PEARS

Codling Moth, Green Apple Aphid, White Apple Leafhopper, Apple Mealybug—12 oz. per 75 gals. dilute spray.

Apple Maggot, Plum Curculio, Rosy Apple Aphid, Woolly Apple Aphid, Red-banded and Fruit Tree Leaf Rollers, Eastern Tent Caterpillar, Pear Psylla, Periodical Cicada, Eye-spotted Bud Moth, European Apple Sawfly, Japanese Beetle, Tentiform Leaf Miners, Forbes, San Jose, Pear Leaf and Lecanium Scales, Bagworm, Green Fruitworm, Apple Rust Mite, Pear Leaf Blister Mite, Pear Rust Mite, Tarnished Plant Bug—2 lbs. per 100 gals. dilute spray. Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, European Fruit Lecanium, Japanese Beetle—3 to 4 lbs. per acre. May be applied on day of harvest.

Fruitworms, Fireworms, Leafhoppers, Cutworms, Japanese Beetle—3 to 6 lbs. per acre.

Grape Leaf Folder, Grape Leafhopper, European Fruit Lecanium, Grape Leaf Skeletonizer—2 to 4 lbs. per acre.

Grape Berry Moth, Japanese Beetle, June Beetles, Red-banded Leaf Roller—4 lbs. per acre.

A dilute spray of 200 gals. per acre is suggested. May be applied on day of harvest.

Mexican Bean Beetle—1 lb. per acre. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetle, Japanese Beetle, Velvetbean Caterpillar—2 lbs. per acre.

Corn Earworm, Armyworms, Stink Bugs, Tarnished Plant Bug—2 to 3 lbs. per acre.

Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Flea Beetles, Leafhoppers—2 to 4 lbs. per acre.

For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

Flea Beetle, Harlequin Bug—1 to 2 lbs. per acre. Corn Earworm, Armyworms, Cabbage Looper—2 to 4 lbs. per acre. (Where cabbage looper is the principal problem use another pesticide. For small loopers after edible portions form, use SEVIN every 5 to 7 days.)

Flea Beetles, Harlequin Bug, Leafhoppers—1 to 2 lbs. per acre.

Six-spotted Leafhopper (Aster Yellows Vector) use 2 to 3 lbs. per acre. Armyworms, Cabbage Looper, Corn Earworm, Tarnished Plant Bug, Stink Bugs—2 to 4 lbs. per acre. Where Cabbage Looper is the principal problem, use an alternative pesticide that is specifically recommended.

Colorado Potato Beetle, Flea Beetles, Leafhoppers—1 to 2 lbs. per acre.

Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs and Lygus Bugs—2 to 4 lbs. per acre.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by burning. Keep out of smoke.

\*Trade-mark of Union Carbide Corporation for the active ingredient Carbaryl.

Distributed by CENOL COMPANY

Libertyville, Illinois 60048

EPA REG. NO. 419-162

E.P.A. EST. No 419-IL-1