

STORAGE AND DISPOSAL

Avoid Storage conditions that might alter the composition or usefulness of the pesticide, e.g. temperature extremes, excessive moisture or humidity, heat, sunlight, friction or contact with irritating substances or media. Keep pesticide container closed in a cool, dry, locked cabinet or room.

Pesticide Disposal:

Do not contaminate water, food, or feed by storage and disposal. Pesticide Wastes are toxic. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Metal Containers: Triple Rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Plastic Containers: Triple Rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, if it is allowed by state and local authorities, by burning. If burned, stay out of smoke.

Household Use Pesticide Only:

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse container. If it can be rinsed thoroughly before discarding in trash.

Cenol

CENOX FOOD PLANT FOGGING INSECTICIDE

— WATER BASE —

A Concentrated PYRENONE Micro Emulsion

- Clear • Stable • Leaves No Oily or Objectionable Residue.

Designed for Use in Edible Product Areas of Official Establishments Operating under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in other Food Processing Plants.

Apply with Conventional Mechanical or Compressed Air Equipment designed and adjusted to deliver a true aerosol mist.

ACTIVE INGREDIENTS:

Pyrethrin 0.01%
Pyrethrin 0.01%
Pyrethrin 0.01%

INERT INGREDIENTS:

Pyrethrin 0.01%
Pyrethrin 0.01%
Pyrethrin 0.01%

PYRENONE - Patent American trademark

100-01-
100-01-
100-01-
100-01-
100-01-

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See rear panel for additional precautionary statements.

Manufactured By:
CENOL Company
Barrington, IL 60010

EPA Reg. No. 419-212
EPA Est. No. 419-IL-1

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONARY STATEMENTS

*HAZARDS TO HUMANS
AND DOMESTIC ANIMALS*

CAUTION

Harmful if swallowed. Avoid contamination of feed, feed and utensils. Avoid contamination of feed and utensils by keeping them well covered while spraying the area.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1,000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of **flies**, including fruit flies, mosquitoes, silverfish, flying moths, gnats, crickets, roaches, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese skippers, and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal moths, and grain mites.

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult **Phorids**, **Sciards**, and **Cecidomyiids** in growing houses, apply at a rate of 1 ounce per 1,000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2,000 cubic feet of space.

CAUTION: Harmful if swallowed. Avoid contamination of feed and feedstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label. Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

419-212