

DIRECTIONS FOR USE

FOR THE SICK ROOM: All articles, including bed-
ding, clothing, etc., wash with a solution of three
ounces of disinfectant to 1 gallon of water.

For TOILETS, CLOTHING and LINEN in sick
rooms: Use in the proportion of three ounces of
disinfectant to one gallon of water. Add the water
to the disinfectant and mix well before using.

WATER CLOSETS and URINALS: Use ounces of
disinfectant to a gallon of water. Apply freely to
all surfaces to be disinfected and scrub the ex-
posed surfaces.

FOR GENERAL DISINFECTING

Remove all articles out from the premises and use
a solution of three ounces of disinfectant to 100 parts of
water. This must be done as a disinfectant in most
cases.

FOR THE TYPHOID SICK ROOM: In the disinfecting
process, all articles, including bedding, clothing, and
other articles, and articles in the sanitary
rooms, should be disinfected. The disinfectant should be
used in the proportion of three ounces to each gallon
of water.

The disinfectant should be used in the proportion of
three ounces to each gallon of water. The disinfectant
should be used in the proportion of three ounces to each
gallon of water.

GARIS: Use in the proportion of three ounces of
disinfectant to each gallon of water. The disinfectant
should be used in the proportion of three ounces to each
gallon of water.

USDA Reg. No. 421-9

NET CONTENTS

421-9

LILAC ODOR INDUCIDE

Proved Coefficient at least 5 by A.O.A.C. Method

Active Ingredients: 20

1. Vegetable Oil Soap
2. Beta Terpineol
3. Isopropyl Alcohol
4. Ortho-phenylphenol
5. Chloro-2-phenylphenol

Total Ingredients: 80 - WATER

Preferable and water to the disinfectant

FOR EXTERNAL USE ONLY

CAUTION - KEEP OUT OF REACH OF CHILDREN
(See side panel for further caution)



A Product of

James Varley & Sons, Inc.

ST. LOUIS, MO

DIRECTIONS: FOR USE

Red Mire: Paint or spray the perches, nest
boxes, drop board, floor and inside walls,
poultry houses and coops with full strength
disinfectant, repeating as often as necessary.
Applied in this manner, it is harmless to the
birds but should not be used directly on the
fowl.

This disinfectant can be used anywhere
where clean and sanitary quarters are desired.
A partial list of places using this type of dis-
infectant follows: Hotels, schools, institutions,
office buildings, factories, jails, public build-
ings, railroad terminals, theatres, stores,
sanitariums and many others.

Keep the barns, kennels, pens and sties thor-
oughly clean by scrubbing and spraying
freely with a solution of disinfectant in the
proportion of one cupful to a pail of water.
Scrub all floorings with same solution. This
will reduce the danger of cross infection.

HATCHERIES: Clean wash floors and trays
with a solution of 1 part disinfectant to 10
parts of water and repeat as often as neces-
sary. In all contagious diseases, separate the
healthy fowls from the affected ones, giving
them fresh clean quarters.

CAUTION: Solutions of this disinfectant may
cause skin irritation in some persons who
might be allergic to this type of Compound.
Such persons who might be allergic to this
Compound should avoid getting solution of
this disinfectant on the skin and if accident-
ly spilled thereon remove by washing with
water.