

BETA-TROL

RED LABEL

Deodorant-Germicide Disinfectant-Cleaner



ACCEPTED
JUN 13 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED
UNDER NO. 421-11. SUBJECT
TO ATTACHED COMMENTS.

Fragrant Lilac Scent

Phenol Coef. 3 – A.O.A.C. Method of Test

ACTIVE INGREDIENTS 50%

1. Beta Terpeneol 2. Vegetable Oil Soap 3. Isopropyl Alcohol

INERT INGREDIENT, Water 50%

CAUTION: KEEP OUT OF REACH OF CHILDREN

NOT GENUINE UNLESS PACKED UNDER THIS LABEL

A Product of *James Varley & Sons, Inc.* ST. LOUIS, MO.

BETA-TROL

DIRECTIONS FOR USE

FLOORS: When mopping or scrubbing use 2½ ounces of BETA-TROL to each gallon of water in the bucket.

GENERAL DISINFECTING: Remove all filth and dirt from the premises and use a solution of one part of BETA-TROL to 60 parts of water.

TOILET ROOMS, WATER CLOSETS AND URINALS: Wash the closet and urinal bowls regularly with a solution made of ½ cupful of BETA-TROL in each gallon of water, leaving the solution in the bowl as long as possible. Scrub the floor and seat frequently with this solution.

GARBAGE: To aid in preventing the breeding of flies and certain other vermin, sprinkle with the solution in the proportion of 4 ounces of BETA-TROL to each gallon of water. Rinse the empty can with some of the solution, leave the wash in the can.

In a typhoid sick-room, in the disinfecting of beds, clothes, utensils and rooms, BETA-TROL should be used in the proportion of four ounces in each gallon of water.

The following table will assist in obtaining accurate dilutions:

1¼ ounces with 1 gallon makes 1% solution
2½ ounces with 1 gallon makes 2% solution
4 ounces with 1 gallon makes 3% solution
6½ ounces with 1 gallon makes 5% solution

Always put BETA-TROL in pail first and then add the water.

For External Use Only – Keep Out of the Reach of Children.

Net Contents: 1 Gallon

USDA Reg. No. 421-11

A Product of *James Varley & Sons, Inc.* ST. LOUIS, MO.

H-D-10. DISINFECTANT

Phenol Coefficient A.O.A.C. Method
S. typhosa 10 S. aureus 10



ACTIVE INGREDIENTS: 35%

Isopropyl Alcohol

Sodium 4 and 6-chloro-2-phenylphenate

Sodium dodecylbenzene sulfonate

Ortho-phenylphenol

INERT INGREDIENT: 65%

Water

Preferably add water to the disinfectant.

ACCEPTED
9-14-72
421-13

DIRECTIONS FOR USE

SURGICAL INSTRUMENTS: Use a 2% solution of Disinfectant for disinfecting surgical instruments and operating equipment for ordinary cases (15 minutes). Use a 3% solution for disinfecting surgical instruments and operating equipment used in connection with infectious cases. Before immersion in the disinfecting solution, surgical instruments should be cleaned of all adhering blood and exudate.

For the **SICK ROOM.** All utensils, such as bedpans, urinals, and wash with a solution of one ounce of Disinfectant to 1-gallon of water.

WATER CLOSETS and URINALS. Use 2 ounces of Disinfectant to a gallon of water. Apply freely to all surfaces to be deodorized and flush the drain pipes with the solution.

GENERAL DISINFECTING AND DEODORIZING

Remove all filth and dirt from the premises and use a solution of one part Disinfectant to 200 parts of water. This mixture acts as a disinfectant against many bacteria and

In the typhoid sick-room, in the disinfecting of bed frames, utensils, and germicide should be used in the proportion of 2 ounces to each gallon of water.

WARNING: Harmful if swallowed. Causes skin and eye irritation. In case of contact, flush with plenty of water; for eyes, get medical attention. Avoid contamination of food.

This Disinfectant is highly recommended for general hospital sanitation: floors, walls, and for the cleaning and disinfecting of sick-room utensils, bed pans, etc. It is especially adapted for the disinfection of instruments, gloves, and used for the disinfection and cleaning of toilet lavatories.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.

CAUTION: KEEP OUT OF REACH OF CHILDREN

WARNING:

Net Contents: _____

USDA Rec. No. 421-13