

Net Contents

# HOSPITAL CRESYLIC DISINFECTANT

COMMERCIAL STANDARD 71/41

PHENOL COEFFICIENT 5 by ~~Food~~ <sup>A.O.A.C.</sup> Method

ACTIVE INGREDIENTS	75%
Cresylic Acid	
Soap	
INERT INGREDIENTS	25%
Water	
Glycerine	

## DIRECTIONS FOR USE

Be sure all surfaces are thoroughly wetted with the disinfecting solution to secure adequate germicidal effect.

**Hospitals:** For disinfecting surgical instruments, use a 2% solution or approximately three ounces in each gallon of water. Instruments should be freed of all adhering blood and serous exudate prior to immersion in the disinfecting solution.

**For Disinfecting Bed Utensils, Etc.:** Use same strength solution.

**As an Antiseptic Dressing for Wounds:** For sores, skin abrasions, wounds, etc., use a 2% solution, applying same to the wound with gauze or absorbent cotton saturated in the solution.

Admirably adapted and specially recommended to Veterinarians in tubercular ~~medication~~ work, and hoof and mouth disease.

ACCEPTED  
Sep 23 1906  
UNDER THE  
FUNGICIDE  
FOR ECONOMIC  
UNDER NO.

PHENOL COEFFICIENT 5 by ~~Food~~ <sup>A.U.A.C.</sup> Method

- ACTIVE INGREDIENTS 75%
  - Cresylic Acid
  - Soap
- INERT INGREDIENTS 25%
  - Water
  - Glycerine

L A C C E

Sep 1900

UNDER THE  
FUNGICIDE  
FOR ECONOMIC  
UNDER NO.

**DIRECTIONS FOR USE**

Be sure all surfaces are thoroughly wetted with the disinfecting solution to secure adequate germicidal effect.

**Hospitals:** For disinfecting surgical instruments, use a 2% solution or approximately three ounces in each gallon of water. Instruments should be freed of all adhering blood and serous exudate prior to immersion in the disinfecting solution.

**For Disinfecting Bed Utensils, Etc.:** Use same strength solution.

~~As an Antiseptic Dressing for Wounds: For sores, skin abrasions, wounds, etc., use a 2% solution, applying same to the wound with gauze or absorbent cotton saturated in the solution.~~

Admirably adapted and specially recommended to Veterinarians in tubercular eradication work, and hoof and mouth disease.

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
**POISON - FOR EXTERNAL USE ONLY**  
**POISONOUS IF TAKEN INTERNALLY**



Internal Antidote—Give milk, white of raw eggs, mucilage or cornstarch paste, and follow with an emetic such as a tablespoonful of mustard in a tumbler of warm water. Avoid contact with skin of undiluted disinfectant.

**Warning:** May be Fatal If Swallowed or Absorbed Through Skin! May Produce Severe Burns! Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling.

Manufactured by  
**JAMES VARLEY & SONS, INC., ST. LOUIS, MO.**