

DIRECTIONS FOR USE

For Disinfecting In:

RESTAURANTS, TAVERNS, FOUNTAINS: First wash glasses, dishes, utensils and cooking equipment in a detergent cleaning solution which does not contain soap. Preferably rinse first with clear water, then rinse again for not less than 30 seconds in a solution containing 2 ounces of Quaternary Ammonium Germicide to each 2 gallons of water. Rinse with potable water. Always allow equipment to air dry. Do not towel.

DAIRIES, FOOD PLANTS, PACKING HOUSES AND FOOD LOCKERS: Surfaces to be disinfected should first be thoroughly cleaned with a soapless cleaner. Then use 1 ounce of Quaternary Ammonium Germicide to 1 gallon of water as a spray or rinse for all working surfaces as well as walls, to control bacteria growth. Work tables, meat blocks and other places with which food may come in contact should be rinsed with clear water after disinfecting.

DEODORIZING REFUSE CONTAINERS, LOCKER ROOMS, DAMP BASEMENTS: Use 2 ounces of Quaternary Ammonium Germicide in ½ gallon of water and spray over garbage and around toilet rooms. This will destroy objectionable odors without leaving a "disinfectant" smell behind.

Net Contents:

EPA Reg. No. 421-154

ACC. 11-2-42
UNDER THE FEDERAL
FUNGICIDE AND RO
FOR ECONOMIC
ED UNDER NO.
TO ATTACHED COMMENT S.



QUATERNARY AMMONIUM GERMICIDE

PHENOL COEFFICIENT 10 A.O.A.C.

"The Versatile Disinfectant"

WARNING: KEEP OUT OF REACH OF CHILDREN

(See side panel for further precautions)

ACTIVE INGREDIENTS 5.0%

Methyldodecylbenzyl trimethyl ammonium chloride
Methyldodecylxylylene bis (trimethyl ammonium chloride)

INERT INGREDIENT, Water 95.0%

A Product of *James Varley & Sons, Inc.* ST. LOUIS, MO.

For Use
SICKRO
nium Ge
several t
This will
dishes a
bedpans
Quaterna
ter for fi
cleaned a

BARBER
combs ar
then soak
of Quate
water.

WARNIN
tion. Do
flush tho
medical a
Avoid coi

FIRST A
gelatin s
large qu
physician

RINSE E
WATER