

DIRECTIONS FOR USE

FOR THE SICK ROOM: All utensils, such as bed-pans, urinals, etc., wash with a solution of 3 ounces of disinfectant to $\frac{1}{2}$ gallon of water to clean and sanitize, applying a fresh solution of the same strength as a final rinse to disinfect.

FOR BOILING CLOTHING and LINEN: used in sick-room. Use in the proportion of 3 ounces of disinfectant to 1 gallon of water. Add the water to the disinfectant and mix well before using.

REST ROOMS and URINALS: Use 3 ounces of sanitizer to 1 gallon of water. Apply freely to all surfaces to be sanitized and scrub the exposed areas.

DAIRIES, FOOD PLANTS, PACKING HOUSES and FOOD LOCKERS: Surfaces to be sanitized should first be thoroughly cleaned with a soapless cleaner. Then use 4 ounces of disinfectant to 1 gallon of water as a spray or rinse for all working surfaces as well as walls, etc., to control bacteria, and mold growth. Work tables, meat blocks and other places with which food may come in contact should be rinsed with clear water after sanitizing. Solutions of the same strength may be used as a flank and udder rinse before milking, as well as a rinse for the hands of workers.

SURGICAL INSTRUMENTS: Use a 3% solution of disinfectant for disinfecting surgical instruments and operating equipment for ordinary cases. Immerse for 15 minutes. Rubber goods should be left in the solution for 30 minutes. Use a 5% solution for disinfecting surgical instruments and operating equipment used in connection with infectious cases. Before immersion in the disinfecting solution, surgical instruments should be cleaned of all adhering blood and exudate.

FOR GENERAL DISINFECTING: First clean to remove all filth and dirt from premises and articles and use a solution of 1 part disinfectant to 50 parts of water. This mixture acts as a disinfectant against *Salmonella typhosa*.

CAUTION

Avoid contamination of food. Avoid getting the concentrate in eyes. In case of contact, wash thoroughly with water. If irritation persists, get medical attention.

LEMON SCENTED DISINFECTANT

Phenol Coef. 5, A.O.A.C. Method

~~REGISTERED TRADE MARK~~

ACTIVE INGREDIENTS: 13%

Isopropyl Alcohol
Methyldodecylbenzyl trimethyl ammonium chloride
Methyldodecylxylene bis—(trimethyl ammonium chloride)
Lemon Essential Oils

INERT INGREDIENT: 87%

Water
Nonylphenoxyethoxyethanol

CAUTION: KEEP OUT OF REACH OF CHILDREN

(See side panel for further cautions)



A Product of

James Varley & Sons, Inc.

ST. LOUIS, MISSOURI

DIRECTIONS FOR USE

IN THE FOOT BATH: A 6% solution of disinfectant is highly recommended for use as a foot bath in locker rooms, gymnasiums, swimming pools, etc. This disinfectant is very effective in destroying the fungus growth causing Athlete's Foot Disease. A 3% solution is also recommended for washing and mopping of floors and benches in locker rooms, swimming pools, gymnasiums, etc., to sanitize and deodorize.

THE TYPHOID SICK ROOM: In the disinfecting of beds, clothing, utensils, typhoid stools, rooms, and the hands of those attending typhoid patients, disinfectant should be used in the proportion of 4 ounces to each gallon of water.

Use $\frac{1}{2}$ cup of the disinfectant to 2 gallons of water for mopping, scrubbing, sprinkling or cleaning purposes. As an antiseptic cleaning solution, use 1 part of disinfectant mixed with 30 parts of water.

This disinfectant is highly recommended for general hospital sanitation; floors, walls, etc., and for the cleaning and disinfecting of sick-room utensils, bed-pans, etc. It is especially adapted for the disinfection of instruments, gloves, and used for the disinfecting and cleaning of toilet lavatories, etc.

This disinfectant can be used anywhere where clean and sanitary quarters are desired. A partial list of places using this type of disinfectant follows: hotels, schools, institutions, office buildings, factories, jails, public buildings, railroad terminals, theatres, stores, sanitariums, and many others.

Every batch is tested for germicidal strength in a modern bacteriological laboratory; but before these tests are made the raw materials are checked in our chemical laboratories—and the finished product is double-checked.

DEODORIZING REFUSE CONTAINERS, LOCKER ROOMS, DAMP BASEMENTS, ETC.: Use 2 ounces of disinfectant in $\frac{1}{2}$ gallon of water and spray over garbage and around toilet rooms, etc. This will destroy objectionable odors without leaving a "disinfectant" smell behind.

USDA Reg. No. 421-251

NET CONTENTS:

VMC

Oct. 26, 1966

DAIRIES, FOOD PLANTS, PACKING HOUSES and FOOD LOCKERS: Surfaces to be sanitized should first be thoroughly cleaned with a soapless cleaner. Then use 4 ounces of disinfectant to 1 gallon of water as a spray or rinse for all working surfaces as well as walls, etc., to control bacteria, and mold growth. Work tables, meat blocks and other places with which food may come in contact should be rinsed with clear water after sanitizing. Solutions of the same strength may be used as a flank and udder rinse before milking, as well as a rinse for the hands of workers. ~~Rinse in POTABLE WATER~~
SURGICAL INSTRUMENTS: Use a 3% solution of disinfectant for disinfecting surgical instruments and operating equipment for ordinary cases. Immerse for 15 minutes. Rubber goods should be left in the solution for 30 minutes. Use a 5% solution for disinfecting surgical instruments and operating equipment used in connection with infectious cases. Before immersion in the disinfecting solution, surgical instruments should be cleaned of all adhering blood and exudate.

FOR GENERAL DISINFECTING: First clean to remove all filth and dirt from premises and articles and use a solution of 1 part disinfectant to 50 parts of water. ~~This mixture acts as a disinfectant against Salmonella typhosa.~~

CAUTION
Avoid contamination of food. Avoid getting the concentrate in eyes. In case of contact, wash thoroughly with water. If irritation persists, get medical attention.

ACTIVE INGREDIENTS: 13%
Isopropyl Alcohol
Methyldodecylbenzyl trimethyl ammonium chloride
Methyldodecylxylylene bis—(trimethyl ammonium chloride)
Lemon Essential Oils
INERT INGREDIENT: 87%
Water
Nonylphenoxypolyethoxyethanol

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See side panel for further cautions)



A Product of

James Varley & Sons, Inc.

ST. LOUIS, MISSOURI

Use 1/2 cup of the disinfectant to 2 gallons of water for mopping, scrubbing, sprinkling or cleaning purposes. As an antiseptic cleaning solution, use 1 part of disinfectant mixed with 30 parts of water.

This disinfectant is highly recommended for general hospital sanitation; floors, walls, etc., and for the cleaning and disinfecting of sick-room utensils, bed-pans, etc. It is especially adapted for the disinfection of instruments, gloves, and used for the disinfecting and cleaning of toilet lavatories, etc.

This disinfectant can be used anywhere where clean and sanitary quarters are desired. A partial list of places using this type of disinfectant follows: hotels, schools, institutions, office buildings, factories, jails, public buildings, railroad terminals, theatres, stores, sanitariums, and many others.


Every batch is tested for germicidal strength in a modern bacteriological laboratory—but before these tests are made the raw materials are checked in our chemical laboratories—and the finished product is then double-checked.

DEODORIZING REFUSE CONTAINERS, LOCKER ROOMS, DAMP BASEMENTS, ETC.: Use 2 ounces of disinfectant in 1/2 gallon of water and spray over garbage and around toilet rooms, etc. This will destroy objectionable odors without leaving a "disinfectant" smell behind.

USDA Reg. No. 421-251

NET CONTENTS:

VMC



DIRECTIONS FOR USE

CONTROLLING THE SPREAD OF ATHLETE'S FOOT FUNGI: A 6% solution is recommended for scrubbing shower room floors, benches, locker room floors, bath mats, swimming pool runways, and other surfaces commonly contacted by the bare feet.

THE TYPHOID SICK ROOM: In the disinfecting of beds, clothing, utensils, rooms and as an aid in the sanitary disposal of typhoid stools, use 4 ounces disinfectant to each gallon of water.