

DURHAM'S

READY TO USE RAT-KIL

READY TO USE PROLIN RAT-KIL WITH WARFARIN*

**WARNING! KEEP OUT OF REACH OF CHILDREN
CHEMISTRY'S GREAT NEW WEAPON
TO FIGHT THE RAT MENACE**

See Antidote and Other Warnings on Back Panel.



**PROLIN RAT-KIL WHEN FED TO RATS
AND MICE ACCORDING TO DIRECTIONS
CAUSES THEM TO DIE OF INTERNAL
HEMORRHAGE.**

PRICE \$1.00

ACTIVE INGREDIENTS:

WARFARIN* (3-p-acetylbenzyl)-4-hydroxycoumarin025%
N-(2-quinoxaliny) sulfanilamide (Sulfaquinoxaline*)025%
INERT INGREDIENTS99.50%

*U. S. PATENT NO. 2,427,578
WISCONSIN ALUMINI RESEARCH FOUNDATION

NET WEIGHT 1 POUND

USDA REGISTRATION NO. 430-5 AUC 2 1966

4 HANDY BAIT CUPS INCLUDED

FOR MICE: Use the 4 handy bait cartons inside this carton.
FOR RATS: We recommend that you remove metal plug from end of this carton and place the open carton where rats can get to it as it makes a good feeding station. CAUTION: Place the cartons in areas inaccessible to children, pets and animals.

WITH WARFARIN-PLUS "PROLIN" DESTROYS HARD-TO-KILL RODENT PESTS

Durham's PROLIN RAT-KIL with WARFARIN contains the well known anti-coagulant WARFARIN and also the new Prolin anti-bacterial agent "Sulfaquinoxaline" which retards the production of Vitamin K in a rodent's digestive system and thereby increases the effectiveness of Warfarin's anti-coagulant properties assuring a much more effective Rodent control.

DIRECTIONS FOR USE

MICE: Place one or two tablespoonfuls of Rat-Kil with Prolin in shallow containers like those enclosed in this carton and place the bait where signs of mice have been seen, spacing the bait stations 8 to 12 feet apart. Keep baits out for 12 or 14 days and replace if they become unpalatable from contamination or spoilage.

RATS: Place 6 to 8 ounces of Rat-Kil in a dry container (this carton makes an excellent bait station) next to burrows, in runways or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 14 days or until all signs of feeding have stopped. Replace stale or spoiled bait with fresh bait. Protect from children, pets or livestock by means of bait boxes or cages.

CAUTION

RAT-KIL contains as its active ingredient an anti-coagulant chemical which if taken accidentally by humans, domestic animals, or pets may reduce the clotting ability of the blood and serious hemorrhage may result. In case RAT-KIL is accidentally eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately. Note For Physicians: When a human has been known to have accidentally ingested RAT-KIL, blood transfusions combined with intravenous and oral doses of Vitamin K are indicated as in case of hemorrhage caused by over-dose of Coumatrol.

Application for Re-Registration for ~~proposed~~ Label for Durham's Rat-Kil with Prolin
Submitted by Durham's Drug Products Company, Company

ACCEPTED