DURHAM'S

5% SEVIN* DUST

FOR

[redacted]

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of dust or spray. Do not get in eyes or on skin. Wash thoroughly after handling. Wear clean clothing. Keep out of the reach of children. Avoid storage near feed and feed products. For the protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. During application avoid drift to adjacent feed and forage plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Burn bag immediately when empty and stay out of the smoke.

ACTIVE INGREDIENT

Carbaryl (1-Naphthyl N-Methylcarbamate)	5%
INERT INGREDIENTS	95%

*Trade Mark of Union Carbide Corp. for the active ingredient (1-Naphthyl N-Methylcarbamate)

CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET WEIGHT 16 OZS. (1 POUND)

MANUFACTURED BY
DURHAM'S DRUG PRODUCTS COMPANY
COMANCHE, TEXAS

APR 22, 1971

430-30

DURHAM'S

DIRECTIONS FOR USING DURHAM'S SEVIN 5% DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS. CONSULT YOUR COUNTY EXTENSION SERVICE AGENT IF IN DOUBT AS TO PROPER USAGE.

DIRECTIONS: BEANS, MEXICAN BEAN BEETLE — ¼ lb. of Durham's Sevin 5% Dust per 1000 sq. ft. LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORMS, FLEA BEETLES, JAPANESE BEETLES — ½ lb. per 1000 sq. ft. CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR — ½ to ¾ lb. per 1000 sq. ft. LIMA BEAN POD BORER, LYGUS and STINKBUGS—1 lb. per 1000 sq. ft. May be used up to harvest. Apply thoroughly at first signs of insect infestation. When necessary repeat every 7 to 10 days.

CUCUMBER, SUMMER SQUASH, PICKLEWORM, MELONWORM — ¼ to ½ lb. per 1000 sq. ft. CUCUMBER BEETLES, SQUASH BUGS, FLEA BEETLES — ½ lb. per 1000 sq. ft. May be used up to harvest.

GRAPES: GRAPE LEAF FOLDER — 1st. brood: ½ lb. per 1000 sq. ft. just before first brood larvae emerge from first rolls. 2nd and 3rd Broods — ½ to 1 lb. per 1000 sq. ft. May be applied up to harvest. GRAPE LEAFHOPPER — ½ lb. per 1000 sq. ft. Thorough coverage is essential. This application may be expected to control DDT resistant leafhoppers.

LETTUCE, MUSTARD GREENS, SPINACH, TURNIPS for Flea Beetle, Harlequin bug, Leafhopper, apply ¾ to 1 lb. per 1000 sq. ft. IMPORTANT — Do not apply to head lettuce within 3 days of harvest. Do not apply to leaf lettuce or Spinach, Turnip or Mustard greens within 14 days of harvest.

DOGS, CATS: BROWN DOG TICKS, FLEAS — Dust liberaly over the animal and rub thoroughly into skin. Apply also to the feet, between the toes and on legs. Dust in and around sleeping quarters and other areas where ticks and fleas occur. Apply at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age.

CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS for northern fowl mite and lice, and as a supplement to previous treatment for chicken mites and fleas, dust birds with 1 lb. of 5% dust per 100 birds. As a litter treatment to control northern fowl mite, chicken mite, lice, fleas and bed bugs, treat litter at rate of 1 lb. per 40 square feet.

Caution: Do not repeat applications within 4 weeks. Do not apply to birds within 7 days of slaughter or do not house birds in treated houses within 7 days of slaughter. Do not apply directly to eggs or litter.

USDA REGISTRATION NO. 430-30

MANUFACTURED BY
DURHAM'S DRUG PRODUCTS COMPANY
COMANCHE, TEXAS