

ACCEPTED  
 MAR 17 1977  
 FEDERAL INSECTICIDE  
 DEFENSIVE ACT  
 SUBJECT

**DURHAM'S**

**DURHAM'S**

**FLEA and TICK  
 POWDER**



FOR  
**DOGS &  
 CATS**

Active Ingredients	
Carbaryl (1-Napthyl Methycarbarmate)	4.5%
Napthalene	.5%
Inert Ingredients	95.0%

Trade Name of Division of ...  
**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET WEIGHT 5 OZS. PRICE

**KILLS FLEAS, TICKS, and LICE  
 CHECKS Fungi and Itching due  
 to Flea, Tick and Lice Bites.  
 HELPS CHECK 'Doggy' ODORS**

**DIRECTIONS**

**FLEAS, BROWN DOG TICKS, LICE** on **DOGS** and **CATS** — Dust powder liberally over animal and rub into skin. Apply also to feet, between toes and legs. Dust in and around sleeping quarters and other areas where fleas or ticks occur. Repeat applications at weekly intervals if needed.

DO NOT USE ON KITTENS less than 4 weeks old.

**CAUTION:** Harmful if swallowed or inhaled. Do not take internally. Avoid breathing the dust when you apply powder. Wash hands after using. Avoid contamination of food and water.

USDA REG. NO. 430-33

DURHAM'S DRUGS PRODUCTS CO. COMANCHE, TEXAS

DURHAM'S

DURHAM'S

# 10% SEVIN<sup>(R)</sup> DUST

### ACTIVE INGREDIENT

Carbaryl (1-Naphthyl N-methylcarbamate) .....	10%
INERT INGREDIENTS .....	90%
Total .....	100%

(R) Registered Trademark of Union Carbide Corporation for Carbaryl Insecticide

**CAUTION: Keep Out of Reach of Children.**  
See Other Caution Statements On Side Panel.

HARMFUL IF INHALED OR SWALLOWED.  
DO NOT TAKE INTERNALLY.  
AVOID BREATHING OF DUST.  
SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT.  
WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK. WEAR REGULAR LONG SLEEVED WORK CLOTHES. CHANGE TO CLEAN CLOTHING DAILY.  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS.  
NOTE FOR PHYSICIAN: CARBARYL IS A MODERATE, REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

### NOTICE TO BUYER

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

### DIRECTIONS FOR USE

To be used as a dust without dilution in conventional ground or air dusters. Dusts should be applied when the wind velocity is less than 5 miles per hour. The best hours for dusting are usually between 5:00 P.M. and 9:00 A.M. Apply on foliage when needed unless otherwise noted in special instructions and repeat at 7 to 10 day intervals if necessary.

CROP AND INSECTS	Rate Per Acre	Interval Last Application to Harvest
ASPARAGUS (On Spears)—Asparagus Beetle Do not repeat application within 3 days. (On Ferns & Brush)—Asparagus Beetle, Apache Cicada Post-Harvest application.	10-20 lbs. 20-40 lbs.	1 day

ASPARAGUS

**ACCEPTED**

BEANS—Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms  
Do not feed storage to dairy animals or animals being finished for slaughter.

CUCUMBERS, SQUASH, CANTALOUPE, WATERMELON—Cucumber Beetle, Spotted Cucumber Beetle, Pickleworm, Leafhoppers, Cucumber Beetle, Fleabeetles  
10 lbs. 0 day

POTATOES, TOMATOES, EGGPLANTS, PEPPERS—Tomato Pinworm, Fall Armyworm, Tomato Hornworm, Tobacco Hornworm, European Corn Borer, Colorado Potato Beetle, Tomato Pinworm, Leafhoppers  
10-20 lbs. 0 day

LETTUCE—Flea Beetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebug, Lygus Bugs  
10-20 lbs. 3 days (Head)  
14 day (Leaf)

CARROTS—Flea Beetles, Leafhoppers, Six-spined Legless Bug (Aster Yellow Vector) Armyworms, Spittlebugs, Lygus Bugs and Stink Bugs  
10-15 lbs.  
10-20 lbs.  
No time limitations.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, KOHLRABI—Fleabeetles, Harlequin Bug  
5-10 lbs. 3 days

CORN EARWORM, ARMYWORMS  
10-20 lbs.

OKRA—Corn Earworm, Stink Bugs  
10-20 lbs. 0 day

CRANBERRIES—Leafhoppers, Fruitworms, European Corn Borer, Japanese Beetle  
15-40 lbs. 1 day

STRAWBERRIES—Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil  
10-20 lbs. 1 day

BLUEBERRY—Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, European Fruit Lecanium, Japanese Beetle  
15-20 lbs. 0 day

GRAPES—Grape Leafhoppers, Grape Leaf Flea, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetle, Red-banded Leaf Roller  
10-30 lbs. 0 day

ORNAMENTALS—Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemum, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose, Rose of Sharon, Yew, Zinnia, Aphids, Bark Beetle, Blister Beetle, Boxelder Bug, Fleabeetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips (Exposed)

Apply to completely cover foliage and repeat as necessary.

E.P.A. REG. NO. 430-49  
Net Weight 50 Lbs.

430-49  
REGISTERED  
MAR 22 1973  
UNDER THE FEDERAL INSECTICIDE ACT AND RELATED ACTS FOR ECONOMIC PEST CONTROL  
EPA UNDER NO. 430-49



70442 UNDER FUNGICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 70443 TO A...

70443

