

DURHAM'S

DURHAM'S

# 10% SEVIN<sup>(R)</sup> DUST

**ACTIVE INGREDIENT**

Carbaryl (1-Naphthyl N-methylcarbamate) ..... 10%

**INERT INGREDIENTS** ..... 90%

Total ..... 100%

(R) Registered Trademark of Union Carbide Corporation for Carbaryl Insecticide

**CAUTION: Keep Out of Reach of Children.**  
See Other Caution Statements On Side Panel.

HARMFUL IF INHALED OR SWALLOWED.  
DO NOT TAKE INTERNALLY.  
AVOID BREATHING OF DUST.  
SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT.  
WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK. WEAR REGULAR LONG SLEEVED WORK CLOTHES. CHANGE TO CLEAN CLOTHING DAILY.  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS.  
NOTE FOR PHYSICIAN: CARBARYL IS A MODERATE, REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NOTICE TO BUYER**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

**DIRECTIONS FOR USE**

To be used as a dust without dilution in conventional ground or air dusters. Dusts should be applied when the wind velocity is less than 5 miles per hour. The best hours for dusting are usually between 5:00 P.M. and 9:00 A.M. Apply on foliage when needed unless otherwise noted in special instructions and repeat at 7 to 10 day intervals if necessary.

**CROP AND INSECTS**

	Rate Per Acre	Interval Last Application to Harvest
ASPARAGUS (On Spears)—Asparagus Beetle	10-20 lbs.	1 day
Do not repeat application within 3 days.		
(On Ferns & Brush)—Asparagus Beetle, Apache Cicada	20-40 lbs.	
Post-Harvest application.		

BEANS—Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Fleabeetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms

Do not feed forage to dairy animals or animals being finished for slaughter.

CUCUMBERS, SQUASH, CANTALOUPE, WATERMELONS—Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Cucumber Beetle, Fleabeetles 10 lbs. 0 day

POTATOES, TOMATOES, EGGPLANTS, PEPPERS—Fleabeetles, Tomato Fruit Worm, Fall Armyworms, Tomato Hornworms, Tobacco Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Tomato Fruitworm, Lace Bug 10-20 lbs. 0 day

LETTUCE—Fleabeetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebugs, Lygus Bugs 10-20 lbs. 3 days (Head) 14 day (Leaf)

CARROTS—Flea Beetles, Leafhoppers, Six-spotted Leafhopper (Aster Yellows Vector) Armyworms, Spittlebugs, Lygus Bugs and Stink Bugs 10-15 lbs. 10-20 lbs. No time limitations.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER, KOHLRABI—Fleabeetles, Harlequin Bug 5-10 lbs. 3 days

Corn Earworm, Armyworms 10-20 lbs. 0 day

OKRA—Corn Earworm, Stink Bugs 10-20 lbs. 0 day

CRANBERRIES—Leafhoppers, Fruitworms, Fireworms, Cutworms, Japanese Beetle 15-40 lbs. 1 day

STRAWBERRIES—Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil 10-20 lbs. 1 day

BLUEBERRY—Blueberry Maggot, Cherry Fruitworm, Cranberry Fruitworm, European Fruit Lecanium, Japanese Beetle 15-20 lbs. 0 day

GRAPES—Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-banded Leaf Roller 10-30 lbs. 0 day

ORNAMENTALS—Arbor Vitae, Azalea, Aster, Birch, Boxwood, Carnation, Camellia, Chrysanthemum, Dahlia, Dogwood, Elm, Gladiolus, Holly, Juniper, Lilac, Marigold, Petunia, Rhododendron, Rose, Rose of Sharon, Yew, Zinnia, Aphids, Bagworm, Blister Beetle, Boxelder Bug, Fleabeetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Orange Tortrix, Periodical Cicada, Plant Bug, Tent Caterpillar, Thrips (Exposed)

Apply to completely cover foliage and repeat as necessary.

E.P.A. REG. NO. 430-49

Net Weight 50 Lbs.

ACCEPTED

MAR 22 1973