

DURHAM'S

FLEA, TICK & LICE DIP

Kills Fleas, Ticks, and Lice
On Dogs and Cats

A Highly Versatile Pyrethrin* Concentrate for Dilution With Water.

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
INERT INGREDIENTS:	94.5%
	100.0%

*Equivalent to 4.0% (butylcarbitol)(6-propylpiperonyl) ether and 1.0% related compounds.

PYRETHRIN - Registered Trademark of Fairfield American Corporation

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel for additional precautions.

NET CONTENTS: ONE QUART
EPA Reg. No. 430.53 EPA Est. No. 430-TX-1

DURHAM'S DRUG PRODUCTS COMPANY
Brownwood, Texas 76801

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES:

Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL:

"Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility."

CONTAINER DISPOSAL:

Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER:

Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

EPA 708

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING

Dogs and Cats: Dilute at the rate of 3 ounces in 1 gallon of water. For rapid control of ticks, fleas and lice, sponge or dip solution on entire animal, thoroughly wetting the skin. Avoid contact with pet's face, eyes and genitalia.

Do NOT treat kittens or puppies less than four weeks old. For rapid control of ticks, fleas, and lice with a mist spray: Dilute at the rate of 1 ounce in 9 ounces of water. To insure thorough coverage, part the fur by stroking against the lay of the hair as spray is applied. Spray ticks directly. Apply to moisten the fur lightly.

To Control Insects in Kennels and Animal Quarters:

Dilute at the rate of 2 ounces in 20 ounces of water.

Remove pets before treating the area and ventilate thoroughly before allowing them to re-enter the treated area. For rapid cleanout of fleas and ticks in kennels, apply with a good machine adjusted to deliver coarse droplets. Apply to the point of runoff, with special attention to cracks and crevices and other insect hiding places. Bedding should be removed and sprayed thoroughly.

AM'S

TICK & DIP

icks, and Lice
and Cats

ncentrate for Dilution With Water.

.....	0.5%
.....	5.0%
.....	94.5%
.....	100.0%

ylcarbityl)(6-propylpiperonyl)
pounds.

of Fairfield American Corporation

CH OF CHILDREN
TION

ditional precautions.

S: ONE QUART
EPA Est. No. 430-TX-1

ODUCTS COMPANY
Texas 76801

STORAGE AND DISPOSAL
Do not contaminate water,
food or feed by storage or
disposal.

**PESTICIDE STORAGE AND
SPILL PROCEDURES:**
Store upright at room tem-
perature. Avoid exposure to ex-
treme temperatures. In case of
spill or leakage, soak up with
an absorbent material such as
sand, sawdust, earth, fuller's
earth, etc. Dispose of with
chemical waste.

PESTICIDE DISPOSAL:
"Wastes resulting from the use
of this product may be dispos-
ed of on site or at an approved
waste disposal facility."

CONTAINER DISPOSAL:
Triple rinse (or equivalent) then
offer for recycling or recondi-
tioning, or puncture and dis-
pose of in a sanitary landfill, or
by other approved State and
local procedures.

**CONTAINERS ONE GALLON
AND SMALLER:**
Do not reuse container. Wrap
container in several layers of
newspaper and discard in
trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

SHAKE WELL BEFORE USING

Dogs and Cats: Dilute at the rate of 3 ounces in 1 gallon of
water. For rapid control of ticks, fleas and lice, sponge or dip
solution on entire animal, thoroughly wetting the skin. Avoid
contact with pet's face, eyes and genitalia.
Do NOT treat kittens or puppies less than four weeks old.
For rapid control of ticks, fleas, and lice with a mist spray:
Dilute at the rate of 1 ounce in 9 ounces of water. To insure
thorough coverage, part the fur by stroking against the lay of
the hair as spray is applied. Spray ticks directly. Apply to
moisten the fur lightly.

To Control Insects in Kennels and Animal Quarters:
Dilute at the rate of 2 ounces in 20 ounces of water.
Remove pets before treating the area and ventilate
thoroughly before allowing them to re-enter the treated area.
For rapid cleanout of fleas and ticks in kennels, apply with a
good machine adjusted to deliver coarse droplets. Apply to
the point of runoff, with special attention to cracks and
crevices and other insect hiding places. Bedding should be
removed and sprayed thoroughly.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid
contamination of feed, food
and utensils. Avoid contami-
nation of food and utensils by
keeping them covered while
spraying the area.
Remove pets, birds and cover
fish aquaria before room ap-
plications are made.

ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do
not apply directly to lakes,
streams or ponds. Do not con-
taminate water by cleaning of
equipment or disposal of
wastes. Apply this product on-
ly as specified on this label.

Buyer assumes all risks of use,
storage and handling of this
product not in strict accor-
dance with instructions given
herewith.

