

17 43-56 PW 117

DURHAM'S

GENERAL PURPOSE INSECTICIDE

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	1.0%
INERT INGREDIENTS	98.9%
	100%

*Equivalent to min. 0.8% (butylcarbitol) propyl piperonyl ether and 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for additional cautions

A Concentrated PYRENONE* Micro Emulsion
Clear-Stable-Leave no oily or objectionable residues
Effective, Fast Acting Insecticide

For use in homes, institutions, restaurants
Food Processing Plants, dairies, livestock and outdoors
for mosquito control

For control of fleas and brown dog ticks on premises
and on pets

Apply with conventional mechanical or compressed air
equipment

EPA REG. NO 430-56

EPA EST. NO. 430-TX-1

Net Contents: ONE QUART

Manufactured For
DURHAM'S DRUG PRODUCTS COMPANY
COMANCHE, TEXAS 76442

STORAGE AND DISPOSAL

Do not contaminate water, food
or feed by storage or disposal

PESTICIDE STORAGE AND SPILL PROCEDURES:

Store upright at room temper-
ature. Avoid exposure to ex-
treme temperatures. In case of
spill or leakage, soak up with an
absorbent material such as
sand, sawdust, earth, fuller's
earth, etc.

Dispose of with chemical waste.

PESTICIDE DISPOSAL:

"Wastes resulting from the use
of this product may be disposed
of on site or at an approved
waste disposal facility."

CONTAINER DISPOSAL:

Triple rinse (or equivalent) then
offer for recycling or recon-
ditioning, or puncture and dispose
of in a sanitary landfill, or by
other approved State and local
procedures.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid con-
tamination of feed and food-
stuffs. Remove pets, birds and
cover fish aquaria before space
spraying or surface applica-
tions

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

INDOOR APPLICATION

The use of this product in food processing establishments should be restricted to those
periods when the plant is not in operation. Food should be removed or covered during treat-
ment. All food processing surfaces should be covered during treatment or thoroughly treat-
ed with potable water before using.

Vacate treated areas and ventilate before reoccupying.
For maximum effectiveness, a combination of spot, surface treatment and fogging is recom-
mended.

SURFACE SPRAY: To control cockroaches, palmto bugs, ants, silverfish spiders,
crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles,
rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore
beetles, meal worms, grain mites and caddises. Use a spray sprayer adjusted to a coarse mist
spray. Direct the spray into hiding places, cracks and crevices, under panels, and on con-
tainers of stored foods, around the base of machinery, behind shelves, and drawers. For
silverfish, spray bookcases. For ants, spray stairs, nests and points of entry. If a treat-
ment only is to be used, spray floors and other surfaces applying a rate of 1 gal. into 750
square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make
localized applications to toys and baseboards. Spray directly into cracks, crevices and in-
fested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take
beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treat-
ment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting
quarters, nearby cracks and crevices along and behind baseboards, mouldings, window
and door frames, and entire areas of floor and floor covering. Fresh bedding should be plac-
ed in animal quarters following treatment. Repeat as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and
air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets
bureaus and other storages directing the spray into cracks, joints and crevices. After a ring
infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid
staining.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a
good machine adjusted to deliver a fine mist or fog. For rapid control of house flies, fruit flies,
gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean
flour moths, Indian meal moths, and tobacco moths: Direct spray at an upward angle
distributing it uniformly throughout the entire area at a rate of 3.6 ounces per 1000 cubic
feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a
rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever
possible.

ON LIVESTOCK: To protect cattle and horses from horn flies, house flies, mosquitoes and
gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse
flies and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the
hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protec-
tion. To control blood sucking lice, apply to the infested areas of the animal, using a stiff
brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

ON PETS: Start spraying at the tail, moving the dispenser rapidly and making sure that the
animal's entire body is covered, including the legs and underbody. While spraying, pull the
hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Do not
spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

OUTDOOR APPLICATION

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For
best results, treat when the wind velocity does not exceed 5 miles per hour. Treat sheltered
or vegetation where mosquitoes may rest, flow, marshy places, stagnant pools or ponds and
shore lines. Apply at a rate of 2.5 pints per acre @ 1 ounce per 1000 square feet.
Using truck mounted conventional misting equipment such as Rotomist® models SIC 91-
100 MT traveling at 5 m.p.h. to treat a 300 foot swath carbide equipment to deliver 0.9
gallon per minute.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or
two glasses of water and induce
vomiting by touching the back of
the throat with finger. Repeat un-
til vomit fluid is clear. Call a
physician immediately. Do not
induce vomiting or give anything
by mouth to an unconscious per-
son.

IF INHALED: Remove victim to
fresh air. Apply artificial res-
piration if indicated.

IF ON SKIN: Remove con-
taminated clothing and wash af-
fected areas with soap and
water.

IF IN EYES: Flush eyes with
plenty of water. Call a physician
if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do
not apply directly to lakes,
ponds, streams, tidal marshes or
estuaries. Do not contaminate
water by cleaning of equipment
or disposal of wastes.

Buyer assumes all risks of use,
storage or handling of this pro-
duct not in strict accordance
with directions given herewith.

ACCEPTED

FEB 28 1986

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under 430-56
EPA Reg. No.